



ISSN Print: 2394-7500
 ISSN Online: 2394-5869
 Impact Factor: 5.2
 IJAR 2015; 1(10): 860-862
 www.allresearchjournal.com
 Received: 05-07-2015
 Accepted: 07-08-2015

R Srivastava
 Department of Mathematics &
 Astronomy Lucknow
 University, Lucknow (India)

An interpolation process with spline of degree six

R Srivastava

Abstract

In this paper, we consider a new technique of spline methods is used for (0, 4, 5) of degree six-lacunary interpolation by splines with functions belonging to $C^{(6)}(I)$ and error bonds, using piecewise polynomials with certain specific properties. Our methods shows better convergence property than the earlier investigations.

Keywords: Lacunary interpolation, quintic splines piecewise polynomial, Spline function.

1. Introduction

Th Fawzy^[3, 4] constructed special kinds of lacunary quintic g-splines and proved that for functions $f \in C^{(4)}$ the method converges faster that investigated by A.K. Verma^[1] and for functions $f \in C^{(5)}$ the order of approximation is the same as the best order of approximation using quintic g-splines. Saxena and Tripathi^[7] have studied splines methods for solving the (0,1,3) interpolation problem. They have used spline interpolants of degree six for functions $f \in C^{(6)}$ to solved the problem. R.S.Misra and K.K. Mathur^[2] solved lacunary interpolation by splines (0;0 2,3) and(0;0,2,4) cases. During the past twentieth both theories of splines and experiences with their use in numerical analysis have undergone a considerable degree of development. According to Fawzy^[3] the interest in spline function is due to the fact that spline function are a good tool for the numerical approximation of functions.

Spline interpolation method, as applied to the solution of differential equation employ some from approximating function such as polynomials to approximate the solution by evaluating the function for sufficient number of points in the domain of the solution. Spline functions are a good tool for the numerical approximation of functions on one hand and they suggest new challenging and rewarding problem's on the other hand. Piecewise linear functions as well as step functions have along been important theoretical and practical tools for approximation of function. Lacunary interpolation by splines appears whenever observation gives scattered or irregular information about a function and its derivatives. The data in the problem of lacunary interpolation has also values of the functions and its derivatives but without Hermite conditions that only consecutive derivative is used at each node. Spline function are arise in many problems of mathematical Physics such as viscoelasticity, hydrodynamics, electromagnetic theory, mixed boundary problems in mathematical physics, biology and Engineering.

The collection of polynomials that form the curve of polynomials that form the curve is collectively referred to as "the spline". The traditional and constrained cubic splines are few different groups of the same family. The group of traditional cubic splines can furthermore be divided into sub group natural, parabolic, runout, cubic run-out and damped cubic splines. The natural cubic spline is by far the most popular and widely used version of the cubic splines family. Spline functions are used in many areas such as interpolation, datafitting, numerical solution of ordinary partial differential equation and also numerical solution of integral equations Lacunary interpolation by splines appears function about a function and its derivatives but without Hermite condition in which consecutive derivatives are used at each nodes, Several researchers have studied the use of spline to solve such interpolation^[5, 8, 9,10, 11] One uses polynomial for approximation because they can be evaluated. cubic spline interpolation is the most common piecewise polynomial method and is referred as "piecewise" since a unique polynomial is fitted between each pair of data points. In addition

Correspondence
R Srivastava
 Department of Mathematics &
 Astronomy Lucknow
 University, Lucknow (India)

the paper mentioned above dealing with best interpolation on approximation by splines there were also few papers that deal with constructive properties of space of splines interpolation. In my earlier work [6] [12] [13] [14] some kinds of lacunary interpolation by g-splines have been investigated. In this paper we will continue to discuss the problem.

This paper is organized as follows- In Section 2, we construct a spline function of degree six which interpolates the lacunary data (0, 4, 5) In section 3 we establish the Error bonds for interpolatory polynomials for $f \in C^{(6)}$ here we also define a Lemma and theorems about spline functions, by using some specific conditions, the method converges faster than the earlier investigations.

2. Spline Interpolant (0, 4, 5) For $f \in C^{(6)}$ (I)

Let $S_{2,\Delta}$ be a piecewise polynomial of degree ≤ 6 The Spline interpolant (0, 4, 5) for function $f \in C^{(6)}$ [I] is given by:

$$(2.1) \quad S_{2,\Delta}(x) = S_{2,k}(x) = \sum_{j=0}^6 \frac{S_{k,j}^{(2)}}{j!} (x - x_k)^j, \quad x_k \leq x \leq x_{k+1}, \quad k = 0(1)n$$

Where $S_{k,j}^{(2)}$, s are explicitly given below in terms of the prescribed data

$$\{f_k^{(j)}\}, j = 0,4,5; k = 0(1)n.$$

For $k = 0(1)n-1$

$$(2.2) \quad S_{k,j}^{(2)} = f_k^{(j)}, j = 0,4,5$$

for $j = 1,2, 3$ and 6 we have

$$(2.3) \quad S_{k,3}^{(2)} = \frac{12}{h^3} [\{f_{x_{k+1}} - f_{x_k} - \frac{h^4}{4!} f_{x_k}^{(4)} - \frac{h^5}{5!} f_{x_k}^{(5)}\} - h \{f_{x_{k+1}}^{(1)} - \frac{h^3}{3!} f_{x_k}^{(4)} - \frac{h^4}{4!} f_{x_k}^{(5)}\} + \frac{h^2}{2} \{f_{x_{k+1}}^{(2)} - \frac{h^2}{2!} f_{x_k}^{(4)} - \frac{h^3}{3!} f_{x_k}^{(5)}\} - \frac{h^3}{12} \{f_{x_{k+1}}^{(3)} - h f_{x_k}^{(4)} - \frac{h^2}{2!} f_{x_k}^{(5)}\}]$$

$$(2.4) \quad S_{k,6}^{(2)} = \frac{72}{h^6} [h \{f_{x_{k+1}}^{(1)} - \frac{h^3}{3!} f_{x_k}^{(4)} - \frac{h^4}{4!} f_{x_k}^{(5)}\} - \{f_{x_{k+1}} - f_{x_k} - \frac{h^4}{4!} f_{x_k}^{(4)} - \frac{h^5}{5!} f_{x_k}^{(5)}\} - \frac{h^2}{2} \{f_{x_{k+1}}^{(2)} - \frac{h^2}{2!} f_{x_k}^{(4)} - \frac{h^3}{3!} f_{x_k}^{(5)}\} + \frac{h^3}{6} \{f_{x_{k+1}}^{(3)} - h f_{x_k}^{(4)} - \frac{h^2}{2!} f_{x_k}^{(5)}\}]$$

$$(2.5) \quad S_{k,2}^{(2)} = \frac{2}{h^2} [\{f_{x_{k+1}} - f_{x_k} - \frac{h^4}{4!} f_{x_k}^{(4)} - \frac{h^5}{5!} f_{x_k}^{(5)}\} - h \{f_{x_{k+1}}^{(1)} - \frac{h^3}{3!} f_{x_k}^{(4)} - \frac{h^4}{4!} f_{x_k}^{(5)}\} + h^2 \{f_{x_{k+1}}^{(2)} - \frac{h^2}{2!} f_{x_k}^{(4)} - \frac{h^3}{3!} f_{x_k}^{(5)}\} - \frac{2}{3} h^3 \{f_{x_{k+1}}^{(3)} - h f_{x_k}^{(4)} - \frac{h^2}{2!} f_{x_k}^{(5)}\} + \frac{55}{6!} h^6 S_{k,6}^{(2)}]$$

$$(2.6) \quad S_{k,1}^{(2)} = \frac{1}{h} [f_{x_{k+1}} - f_{x_k} - \frac{h^4}{4!} f_{x_k}^{(4)} - \frac{h^5}{5!} f_{x_k}^{(5)} - \frac{h^2}{2!} S_{k,2}^{(2)} - \frac{h^3}{3!} S_{k,3}^{(2)} - \frac{h^6}{6!} S_{k,6}^{(2)}]$$

3. Error Bond for Spline Interpolants.

Suppose $f \in C^{(6)}$ [II], then by the Taylor expansions, we establish the following Lemma by using the modulus of continuity $\omega(f^{(6)}; h)$.

Lemma 3.1

For $j = 1, 2, 3$ and 6 we have

$$|S_{k,j}^{(2)} - f_k^{(j)}| \leq C_{k,j}^{(2)} h^{6-j} \omega(f^{(6)}; h),$$

$J = 1, 2, 3$ and 6 .

$K = 0(1)n-1$

Where the constants $C_{k,j}^{(2)}$ are given by:

$$C_{k,1}^{(2)} = \frac{83}{3600}, \quad C_{k,2}^{(2)} = \frac{29}{60}, \quad C_{k,3}^{(2)} = \frac{4}{15}, \quad C_{k,6}^{(2)} = \frac{13}{5}$$

Proof.

For $j = 1, 2, 3$ and 6 Using Taylor's expansion from (2.1)-(2.6), we have

$$(3.1) \quad |S_{k,1}^{(2)} - f_k^{(1)}| \leq \frac{83}{3600} h^5 \omega(f^{(6)}; h),$$

$$(3.2) \quad |S_{k,2}^{(2)} - f_k^{(2)}| \leq \frac{29}{60} h^4 \omega(f^{(6)}; h),$$

$$(3.3) \quad |S_{k,3}^{(2)} - f_k^{(3)}| \leq \frac{4}{15} h^3 \omega(f^{(6)}; h), \text{ and}$$

$$(3.4) \quad |S_{k,6}^{(2)} - f_k^{(6)}| \leq \frac{13}{5} \omega(f^{(6)}; h),$$

This completes the Proof of the Lemma 3.1

Theorem 3.2

Let $f \in C^{(5)}$ (I) and $S_{2,\Delta} \in C^{(0,4,5)}$ [I] be the unique spline interpolant (0, 4, 5) given in (2.1) - (2.6), then

$$(3.5) \quad \| D^{(j)} (f - S_{2,\Delta}) \|_{L_\infty [x_k, x_{k+1}]} \leq c_{2,k}^j h^{6-j} \omega(f^{(6)}, h), \quad j=0(1)6; \quad k=0(1)n-1$$

Where the constants $c_{2,k}^{(j)}$'s are given by:

$$c_{2,k}^{(0)} = \frac{6557}{450}, \quad c_{2,k}^{(1)} = \frac{2381}{3600}, \quad c_{2,k}^{(2)} = \frac{103}{120}, \quad c_{2,k}^{(3)} = \frac{21}{30}, \\ c_{2,k}^{(4)} = \frac{13}{10}, \quad c_{2,k}^{(5)} = \frac{13}{5}, \quad c_{2,k}^{(6)} = \frac{13}{5}$$

Proof:

For $k = 0(1)n-1, j = 0(1)6$

$$|f(x) - S_{2,\Delta}| \leq |f(x) - S_k(x)|$$

$$\leq \sum_{j=0}^5 \frac{|f^{(j)}(x_k) - s_k^{(j)}| h^j}{j!} + \frac{|f^{(6)}(\delta_k) - s_k^{(6)}| h^6}{6!}$$

Where $x_k < S_k < x_{k+1}$ Using Lemma 3.1 and the definition of the modulus of continuity of $f^{(6)}(x)$, we obtain

$$(3.6) \quad |f(x) - S_k(x)| \leq \frac{6557}{450} h^6 \omega(f^{(6)}; \delta),$$

$$(3.7) \quad |f_{x_k}^{(1)} - S_k^{(1)}(x)| \leq \frac{2381}{3600} h^5 \omega(f^{(6)}; h),$$

$$(3.8) \quad |f_{x_k}^{(2)} - S_k^{(2)}(x)| \leq \frac{103}{120} h^4 \omega(f^{(6)}; h),$$

$$(3.9) \quad |f_{x_k}^{(3)} - S_k^{(3)}(x)| \leq \frac{21}{30} h^3 \omega(f^{(6)}; h),$$

$$(3.10) \quad |f_{x_k}^{(4)} - S_k^{(4)}(x)| \leq \frac{13}{10} h^2 \omega(f^{(6)}; h),$$

$$(3.11) \quad |f_{x_k}^{(5)} - S_k^{(5)}(x)| \leq \frac{13}{5} h \omega(f^{(6)}; h), \text{ and}$$

$$(3.12) \quad |f_{x_k}^{(6)} - S_k^{(6)}(x)| \leq \frac{13}{5} \omega(f^{(6)}; h),$$

Using (3.6)-(3.12) this completes the Proof of the Theorem 3.1.

4. Conclusion

In this paper, we have studied the existence and uniqueness of (0, 4, 5) of degree six and error bonds for interpolants for functions belonging to $C^{(6)}$ (I). Our methods are having better convergence property Also we conclude that this new

technique we used in proving of the Lemma and one important theorem of spline function is far more better than the earlier investigations.

5. References

1. Varma AK. Lacunary interpolation by splines-II Acta Math. Acad. Sci. Hungar 1978; 31:193-203.
2. Misra RS, Mathur KK. Lacunary interpolation by splines (0; 0, 2, 3) and (0; 0, 2, 4) cases, Acta Math. Acad. Sci. Hungar 1980; 36(3-4):251-260.
3. Fawzy Th. Notes on Lacunary interpolation with splines-III, (0, 2) – interpolation with quintic g-splines Acta Math. Hung 1987; 50(1-2):33-37.
4. Fawzy Th. (0, 1, 3) Lacunary interpolation by G-splines, Annales Univ. Sci., Budapest, Section Maths. XXXIX, 1986, 63-67.
5. Györvari J. Lacunary interpolation spline functionen, Acta Math. Acad. Sci. Hungar 1983; 42(1-2):25-33.
6. Srivastava R. lacunary interpolation by g-splines: International journal of Mathematics and Computer Research, 2014, 2(12).
7. Saxena RB. Tripathi HC. (0, 2, 3) and (0, 1, 3)-interpolation by six degree splines, Jour. Of computational and applied Maths, 1987; 18:395-101.
8. Pandey Ambrish Kumar, Ahmad QS. Singh Kulbhushan: Lacunary Interpolation (0, 2; 3) problem and some comparison from Quartic splines: American journal of Applied Mathematics and statistics. 2013; 1(6):117-120.
9. Lang F, Xu X. “A new cubic B-spline method for linear fifth order boundary value problems”. Journal of Applied Mathematics and computing, 2011; 36:1-2:110-116.
10. Abbas Y, Albayati Rostam KS, Faraidun K Hamasalh. Consturction of Lacunary Sixtic spline function Interpolation and their Applications. Mosul University, J Edu and Sci. 2010, 23(3).
11. Jwamer KH. Radha GK. Generalization of (0, 4) lacunary interpolation by quantic spline. J. of Mathematics and Statistics; New York 2010; 6(1):72-78.
12. Srivastava R. On lacunary Interpolation through g-splines: International journal of Innovative Research in Science, Engineering and Technology, 2015, 4(6).
13. Srivastava R. A new kind of Lacunary Interpolation through g-Splines: International journal of Innovative Research in Science, Engineering and Technology, 2015, 4(8).
14. Srivastava R. A Lacunary Interpolation with splines of degree six International Journal of Recent Scientific Research Research 2015; 6(8):5824-5826.