



ISSN Print: 2394-7500
ISSN Online: 2394-5869
Impact Factor: 5.2
IJAR 2016; 2(9): 94-96
www.allresearchjournal.com
Received: 14-07-2016
Accepted: 15-08-2016

Katari Kantha
Department of Community
Health Nursing, Narayana
College of Nursing, Nellore,
Andhra Pradesh, India

Arumugam Indira
Principal, Narayana College of
Nursing, Nellore, Andhra
Pradesh, India

Prevalence of hypertension among the adults in Koratur VS Kovur Nellore

Katari Kantha and Arumugam Indira

Abstract

Background: Hypertension is the new-era pandemic, which causes about 7.1 million deaths per year globally. Various risk factors are implicated in the development of hypertension and there are differences in these risk factors in coastal and noncoastal populations depending on the level of development and epidemiological transition.

Aim: to assess the prevalence of hypertension

Setting and Design: The study was conducted in Koratur (coastal area) and Kovur (non coastal area) by using a descriptive design.

Materials and Methods: A total of 500 samples were included in this study. Among this, 250 samples belongs to coastal area and 250 samples belongs to non coastal area by using convenience sampling technique.

Statistical Analysis Used: The collected data was organized, tabulated, analyzed and interpreted by using descriptive and inferential statistics based on the objectives of the study.

Results: In Koratur, Out of 250 samples, With regard to the category of the blood pressure 27(10.8%) had stage-I hypertension, 5(2%) had stage-II hypertension, 42(16.8%) had grade-I isolated systolic hypertension, and 7(2.8%) had grade-II isolated systolic hypertension. Known Hypertensive cases are 38(15.2%), Newly diagnosed cases are 43(17.2%). With regard to BMI, among 250 samples 22(8.8%) were overweight and 15(6%) were obese. in Kovur, among 250 samples, 47(18.8%) had stage-I hypertension, 13(5.2%) had stage-II hypertension, 2(0.8%) had stage-III hypertension, 47(18.8%) had grade-I hypertension, and 2(0.8%) had grade-II hypertension. Known Hypertensive cases are 9(3.6%). Newly diagnosed cases are 102(40.8%). With regard to BMI among 250 samples 37(14.8%) were overweight and 9(3.6%) were obese.

Conclusion: The above results shown that blood pressure values are high in the Kovur (non coastal area) than Koratur (coastal area).

Keywords: Hypertension, non coastal area, coastal area, heart attack, stroke

1. Introduction

One in three adults worldwide has high blood pressure. Hypertension increases the risk of heart attack, stroke, kidney failure and much other associated co morbidity. Treating raised blood pressure and maintaining it below 140/90 mmHg is associated with a reduction in cardiovascular complication. The theme for World Health Day (WHD) 2013 is "high blood pressure". The goal of WHD 2013 is to reduce heart attacks and strokes. Keeping in line with the WHO, Government of India, Country Cooperation Strategy, the WHO 2013 events in India are aimed at raising the awareness amongst national policymakers, program managers and other stakeholders on the need to strengthen the Indian health system to make it competent enough to respond to hypertension and related co morbidities [1].

Kantha, K and Indira, A. (2015) conducted a cross sectional study on prevalence of hypertension among the adults in coastal and non coastal areas. A total of 5000 samples were included in the study. In that 2500 samples belongs to coastal areas and 2500 samples belongs to non coastal areas. The prevalence of stage-I hypertension in coastal areas is 460(18.4%) but in non coastal areas it is 1413(56.50%). The results indicate that there is high prevalence of hypertension in non coastal areas than coastal areas [2].

Correspondence
Katari Kantha
Department of Community
Health Nursing, Narayana
College of Nursing, Nellore,
Andhra Pradesh, India

Arumugam Indira *et al.* (2015) conducted a study on prevalence of prehypertension among the adults in coastal and non coastal areas. The study results shown that regarding prehypertension in SBP, in coastal areas 1129(45.16%) and in non coastal areas 971(38.84%). The results indicate that there is high prevalence of pre hypertension in coastal areas than non coastal areas. Further studies are needed to find out the reasons and measures to control high blood pressure is necessary [3].

Even today there is scarcity of the studies in coastal and non coastal areas of India. With this background, present study has been undertaken to study the prevalence of hypertension.

2. Objectives of the Study

- To assess the prevalence of hypertension among adults of coastal and non coastal areas.
- To identify the risk factors of hypertension among adults of coastal and non coastal areas.
- To compare the prevalence of hypertension between coastal and non coastal areas.
- To find association between the prevalence of hypertension with selected socio demographic variables.

3. Detailed Research Plan

3.1 Research Approach: Quantitative Approach.

3.2 Research Design: Descriptive design.

3.3 Research Setting: The study was conducted in Koratur (coastal area) and Kovur (non coastal area) by using a descriptive design.

3.4 Coastal area means areas within 2km from mean low water mark (MLWM) or mean high water mark (MHWM).

3.5 Non-coastal area means areas far 2km from mean low water mark (MLWM) or mean high water mark (MHWM).

3.6 Sampling Technique: Convenience sampling technique

3.7 Sample Size: A total of 500 samples were included in this study. Among this, 250 samples belongs to Koratur (coastal area) and 250 samples belongs to Kovur (non coastal area).

4. Results and discussion

4.1 Comparison of Blood Pressure in Koratur and Kovur.

Table 1: Comparison of Blood Pressure in Koratur and Kovur. (N=250)

| Blood Pressure Category | Koratur | | Kovur | | Correlation coefficient | Standard deviation |
|-------------------------|---------|-------|-------|-------|-------------------------|--------------------|
| | (f) | (%) | (f) | (%) | | |
| Optimal | 55 | 22% | 14 | 5.6% | 0.67 | 25.17 |
| Norma | 65 | 26% | 50 | 20% | | |
| High Normal | 49 | 19.6% | 75 | 30% | | |
| Stage-I | 27 | 10.8% | 47 | 18.8% | | |
| Stage-II | 5 | 2% | 13 | 5.2% | | |
| Stage-III | 0 | 0.00% | 2 | 0.8% | | |
| Grade-I | 42 | 16.8% | 47 | 18.8% | | |
| Grade-II | 7 | 2.8% | 2 | 0.8% | | |




Fig 1: Comparison of Blood Pressure in Koratur and Kovur

The prevalence of stage-I BP in coastal area is 27(10.8%) but in non coastal areas it is 47(18.8%). The correlation

coefficient value is highly significant (0.67) and the standard deviation is 25.17.

4.2 Comparison of Body Mass Index in Koratur and Kovur




Fig 2: Comparison of Body Mass Index in Koratur and Kovur

The prevalence of overweight samples in coastal area is 22 (8.8%), obesity is 15(6%) but in non coastal areas it is 37(14.8%) and 9(3.6%). The correlation coefficient value is highly significant (0.99) and the standard deviation is 88.97.

4.3 Association of Socio Demographic Data with the Blood Pressure in Koratur: There is a significant association of demographic variables with age, family, Working members in family, Type of salt used, amount of vegetables per day, habits, worship of god, are you a known hypertensive, entertainment and remaining are non significant.

4.4 Association of Socio Demographic Data with the Blood Pressure in Kovur: There is a significant association of demographic variables with sleeping pattern, exercise, hotel food, worship of god, are you a known hypertensive and remaining are non significant.

5. Conclusion

- The above results shown that stage-1 and stage-2 values are higher in the Kovur (non-coastal area) than in the Koratur (coastal area).
- Among hypertension cases the prevalence of over weight is more in non-coastal area than coastal area.
- The variables like Age, exercise, Type of oil used for cooking, Type of salt used, habits, intake of fish, are you having stress and are you a known hypertensive are the influencing risk factors for the development of hypertension among the adults.

6. References

1. Anchala R, Kannuri NK, Pant H, *et al.* Hypertension in India: a systematic review and meta-analysis of prevalence, awareness, and control of hypertension. *J Hypertens.* 2014; 32(6):1170-7.
2. Katari Kantha, Arumugam Indira. Prevalence of hypertension among the adults in coastal and non coastal areas *International Journal of Development Research.* 2015; 05(01):3134-3139.
3. Arumugam Indira *et al.*, Prevalence of Pre Hypertension among the Adults Aged 20-60 Years in Coastal and Non Coastal areas *International Journal of recent scientific research.* 2015; 6(11):7166-7170.

4. Arumugam Indira, Katari Kantha. Nutritional status of antenatal and postnatal mothers in selected rural areas at Nellore. *International Journal of Recent Scientific Research.* 2015; 6(2):2796-2800.
5. Katari Kantha *et al.*, Knowledge of the Expectant Fathers regarding safe Motherhood in Narayana Medical College Hospital, Nellore. *International Journal of Recent Scientific Research.* 2015; 6(4):3423-3427.
6. Vanaja Kumari B *et al.*, Comparative Study to Assess the Knowledge Regarding Behavioural Problems of Underfive Children Among Employed and Unemployed Mothers. *International Journal of Recent Scientific Research.* 2015; 6(5):4183-4187.
7. Katari Kantha *et al.*, Nutritional Status of Infant Children In Selected Rural Areas At Nellore *International Journal Of Recent Scientific Research.* 2015; 6(6):4597-4601.
8. Katari Kantha, Usha Rani M, Alfia Parameswaran, Arumugam Indira. The knowledge regarding eating disorders among adolescent girls *International Journal of Applied Research.* 2016; 2(5):864-866.
9. Katari Kantha, Arundhathi S, Arumugam Indira, Laleima Chanu. Knowledge regarding management of dengue fever among staff nurses *International Journal of Applied Research.* 2016; 2(6):147-149.
10. Katari K. Prevalence of hypertension among the adults in Indukurpet Vs Vidavalur, Nellore. *NNJ.* 2015; 4(4):30-33.