



ISSN Print: 2394-7500  
 ISSN Online: 2394-5869  
 Impact Factor: 5.2  
 IJAR 2015; 1(12): 524-526  
 www.allresearchjournal.com  
 Received: 08-09-2015  
 Accepted: 10-10-2015

**Parvinder Singh**  
 P. G. Department of  
 Mathematics, S.G.G.S. Khalsa  
 College, Mahilpur.  
 (Hoshiarpur), Punjab, India.

## The Extension of Measures

**Parvinder Singh**

### Abstract:

In this paper we have to show that if  $A$  is any algebra of subsets of a set  $X$  and if  $\mu$  is a measure defined on  $A$  then there exist a  $\sigma$ - algebra  $A^*$  containing  $A$  and a measure  $\mu^*$  defined on  $A^*$  such that  $\mu^*(E) = \mu(E)$  in  $A$ . In other words the measure  $\mu$  can be extended to a measure on a  $\sigma$ -algebra  $A^*$  of subsets of  $X$ , which contains  $A$ .

**Keywords:** Algebra,  $\sigma$ - algebra, Measure, Measurable set, Finite and Infinite Measure.

### 1.1 Introduction:

Let  $X$  be any set and  $\mathcal{A}$  be any non-empty subset of  $P(X)$  the power set of  $X$ ,  $\mathcal{A}$  is said to be an algebra if

1.  $\phi$  and  $X \in \mathcal{A}$
2.  $A, B \in \mathcal{A} \Rightarrow A \cup B \in \mathcal{A}$
3.  $A \in \mathcal{A} \Rightarrow A^c \in \mathcal{A}$

Here it is easy to prove that if  $\mathcal{A}$  is an algebra and  $A$  and  $B$  are in  $\mathcal{A}$  then  $A \cap B$  and  $A/B$  are also in  $\mathcal{A}$  and also  $A \Delta B = (A - B) \cup (B - A)$  belongs to  $\mathcal{A}$ .

### 1.2 Remark:

$\mathcal{A}$  is closed under finite unions and finite intersections, that if  $A_1, A_2, \dots, A_n \in \mathcal{A}$  then

$$\begin{cases} \bigcup_{i=1}^n A_i \in \mathcal{A} \\ \bigcap_{i=1}^n A_i \in \mathcal{A} \end{cases}$$

### Example 1:

If  $X = [0, 1)$ , then the set  $\mathcal{A}$  consisting of  $\phi$  and all unions  $A = \bigcup_{i=1}^n [a_i, b_i)$  where  $0 \leq a_i < b_i \leq a_{i+1} \leq 1$  is algebra.

**Example 2:** If  $X$  be a finite set then consider the set

$E = \{A \in P(X) / A \text{ is finite or } A^c \text{ is finite}\}$  then  $E$  is an algebra. It can be easily verified that  $E$  is closed under finite unions and finite intersections.

**Example 3:** Obviously for any set  $X$  the power set  $P(X)$  and  $E = \{\phi, X\}$  are  $\sigma$ -algebras in  $X$  and  $P(X)$  is called the largest and  $E = \{\phi, X\}$  is the smallest  $\sigma$ -algebra.

**1.3 Definition:** If  $B$  be an arbitrary subset of  $X$ , define  $\mu^*(B) = \text{Inf} \sum_{i=1}^{\infty} \mu(E_i)$  where the infimum is extended over all sequences  $\{E_i\}$  of sets in  $A$  such that  $B \subseteq \bigcup_{i=1}^{\infty} E_i$ . Generally the function  $\mu^*$  defined above is called outer measure generated by  $\mu$ .

**1.4 Lemma:** The function  $\mu^*$  defined above satisfy the following

- (1)  $\mu^*(\phi) = 0$
- (2)  $\mu^*(A) \geq 0$  For any  $A \subseteq X$ .
- (3) If  $A \subseteq B$ , then  $\mu^*(A) \leq \mu^*(B)$ .
- (4) If  $A \in \mathcal{A}$  then  $\mu^*(A) = \mu(A)$ .
- (5) If  $(E_i)$  be a sequence of subsets of  $X$ , then  $\mu^*(\bigcup_{i=1}^{\infty} E_i) \leq \sum_{i=1}^{\infty} \mu^*(E_i)$ .

**Correspondence**  
**Parvinder Singh**  
 P. G. Department of  
 Mathematics, S.G.G.S. Khalsa  
 College, Mahilpur.  
 (Hoshiarpur), Punjab, India.

**Proof:**

Statements (1), (2) and (3) are immediate consequences of the definition 1.1.

(4) Since  $\{A, \phi, \phi, \dots\}$  is a countable collection of sets in  $\mathcal{A}$  whose union contains A, then it follows that  $\mu^*(A) \leq \mu(A) + 0 + 0 + \dots = \mu(A)$ ..... (1)

Conversely, If  $(E_i)$  be any sequence of subsets from  $\mathcal{A}$  with  $A \subseteq \cup E_i$ , then

$A = \cup (A \cap E_i)$ . Since  $\mu$  is a measure on  $\mathcal{A}$ , then  $\mu(A) \leq \sum_{i=1}^{\infty} \mu(A \cap E_i) \leq \sum_{i=1}^{\infty} \mu(E_i)$ .

Follows that  $\mu(A) \leq \mu^*(A)$ .....(2)

Hence from (1) and (2) we have  $\mu^*(A) = \mu(A)$ .

(5) Let  $\varepsilon > 0$  be arbitrary and for each n chose a sequence  $(E_{nk})$  of the sets in  $\mathcal{A}$  such that  $B_n \subseteq \cup_{k=1}^{\infty} E_{nk}$  and  $\sum_{k=1}^{\infty} \mu(E_{nk}) \leq \mu^*(B_n) + \frac{\varepsilon}{2^n}$ . Since  $\{E_{nk} : n, k \in \mathbb{N}\}$  is a countable collection from  $\mathcal{A}$  whose union contains  $\cup B_n$  it follows from the definition of  $\mu^*$  that  $\mu^*(\cup_{i=1}^n B_n) \leq \sum_{n=1}^{\infty} \sum_{k=1}^{\infty} \mu(E_{nk}) \leq \sum_{n=1}^{\infty} \mu^*(B_n) + \varepsilon$ . Since  $\varepsilon$  is arbitrary, hence proved.

**1.5 Remark:**

The property (5) is referred by saying that  $\mu^*$  in countably sub-additive. Although  $\mu^*$  has the advantage that it is defined for arbitrary subsets of X, it has the defect that it is not necessarily countably additive. Hence we are willing to restrict  $\mu^*$  to a smaller  $\sigma$ -algebra, provided we can find one containing  $\mathcal{A}$  and over which  $\mu^*$  has the property of countable additivity. There is a remarkable condition due to Caratheodory which provides the desired restriction of the domain of  $\mu^*$ .

**1.6 Definition:**

A subset E of X is said to be  $\mu^*$ -measurable if

$$\mu^*(A) = \mu^*(A \cap E) + \mu^*(A \setminus E) \dots \dots \dots (A)$$

For all subsets A of X. The collection of all  $\mu^*$ -measurable sets is denoted by  $\mathcal{A}^*$ . Condition (A) indicated an additivity property on  $\mu^*$

**1.7 Caratheodory Extension theorem:**

The collection  $\mathcal{A}^*$  of all  $\mu^*$ -measurable sets is an  $\sigma$ -algebra containing  $\mathcal{A}$ . Moreover if  $(E_n)$  is a disjoint sequence in  $\mathcal{A}^*$ , then

$$\mu^*(\cup_{n=1}^{\infty} E_n) = \sum_{n=1}^{\infty} \mu^*(E_n).$$

**Proof:**

It is clear that  $\phi$  and X are  $\mu^*$ -measurable sets and if  $E \in \mathcal{A}^*$  then its complement  $X \setminus E$  belongs to  $\mathcal{A}^*$ . Next we shall show that  $\mathcal{A}^*$  is closed under intersections. Indeed suppose that E and F are  $\mu^*$ -measurable. Then for any  $A \subseteq X$  and  $E \in \mathcal{A}^*$ , we have

$$\mu^*(A \cap F) = \mu^*(A \cap F \cap E) + \mu^*((A \cap F) \setminus E)$$

$$\text{Since } F \in \mathcal{A}^* \text{ then } \mu^*(A) = \mu^*(A \cap F) + \mu^*(A \setminus F).$$

Let  $B = A \setminus (E \cap F)$ , then it is readily seen that  $B \cap F = A \cap F \setminus E$  and  $B \setminus F = A \setminus F$ ; since  $F \in \mathcal{A}^*$  it follows that

$$\mu^*((A \cap F) \setminus E) = \mu^*((A \cap F) \setminus E) + \mu^*(A \setminus F)$$

Combining these three relations we get

$$\mu^*(A) = \mu^*(A \cap E \cap F) + \mu^*(A \setminus (E \cap F)).$$

Which shows that  $E \cap F$  belongs to  $\mathcal{A}^*$ . Since  $\mathcal{A}^*$  is closed under intersection and complementation, it follows that  $\mathcal{A}^*$  is an algebra.

Suppose that  $E, F \in \mathcal{A}^*$  and  $E \cap F = \emptyset$ . If we take A to be  $A \cap (E \cup F)$  in (A) we get

$$\mu^*(A \cap (E \cup F)) = \mu^*(A \cap E) + \mu^*(A \cap F)$$

For  $A = X$ , this relation implies that  $\mu^*$  is additive on  $\mathcal{A}^*$ .

We shall now show that  $\mathcal{A}^*$  is a  $\sigma$ -algebra and that  $\mu^*$  is countably additive on  $\mathcal{A}^*$ . Let  $(E_i)$  be a disjoint sequence in  $\mathcal{A}^*$  and let  $E = \cup E_k$  if A be any subset of X then

$$\begin{aligned} \mu^*(A) &= \mu^*(A \cap F_n) + \mu^*(A \setminus F_n) \\ &= \sum_{k=1}^{\infty} \mu^*(A \cap E_k) + \mu^*(A \setminus F_n). \end{aligned}$$

Since  $F_n \subseteq E$ , then  $A \setminus E \subseteq A \setminus F_n$  and letting  $n \rightarrow \infty$  the above relation yields

$$\sum_{k=1}^{\infty} \mu^*(A \cap E_k) + \mu^*(A \setminus E) \leq \mu^*(A).$$

On the other hand it follows from Lemma 1.5 (5) that  $\mu^*(A \cap E) \leq \sum_{k=1}^{\infty} \mu^*(A \cap E_k)$ ,  $\mu^*(A) \leq \mu^*(A \cap E) + \mu^*(A \setminus E)$  on combining these inequalities we get that

$$\begin{aligned} \mu^*(A) &= \mu^*(A \cap E) + \mu^*(A \setminus E) \\ &= \sum_{k=1}^{\infty} \mu^*(A \cap E_k) + \mu^*(A \setminus E). \end{aligned}$$

In particular this shows that  $E = \cup_{k=1}^{\infty} E_k$  is

$\mu^*$ -measurable. on taking  $A = E$ , we obtain

$$\mu^*(\cup_{n=1}^{\infty} E_n) = \sum_{n=1}^{\infty} \mu^*(E_n).$$

Now it remains to prove that  $A \subseteq \mathcal{A}^*$ . It was proved in lemma 1.5(4) that if  $E \in \mathcal{A}$ , then  $\mu^*(E) = \mu(E)$ , but we need to show that E is  $\mu^*$ -measurable. Let A be an arbitrary subset of X; it follows from 1.5 (5) that  $\mu^*(A) \leq \mu^*(A \cap E) + \mu^*(A \setminus E)$ . To establish the opposite inequality, let  $\varepsilon > 0$  be arbitrary and let  $(F_n)$  be a sequence in  $\mathcal{A}$  such that

$$A \subseteq \cup F_n \text{ and } \sum_{n=1}^{\infty} \mu(F_n) \leq \mu^*(A) + \varepsilon.$$

Since  $A \cap E \subseteq \cup (F_n \cap E)$  and  $A \setminus E \subseteq \cup (F_n \setminus E)$ , it follows that

$$\mu^*(A \cap E) \leq \sum_{n=1}^{\infty} \mu(F_n \cap E), \quad \mu^*(A \setminus E) \leq \sum_{n=1}^{\infty} \mu(F_n \setminus E).$$

Hence we have

$$\begin{aligned} \mu^*(A \cap E) + \mu^*(A \setminus E) &\leq \sum_{n=1}^{\infty} \{\mu(F_n \cap E) + \mu(F_n \setminus E)\} = \\ &= \sum_{n=1}^{\infty} \mu(F_n) \leq \mu^*(A) + \varepsilon. \end{aligned}$$

Since  $\varepsilon$  is arbitrary, the desired inequality is established and the set E belongs to  $\mathcal{A}^*$ .

The Caratheodory Extension Theorem shows that a measure  $\mu$  on an algebra  $\mathcal{A}$  can always be extended to a measure  $\mu^*$  on a  $\sigma$ -algebra  $\mathcal{A}^*$  containing  $\mathcal{A}$ . The  $\sigma$ -algebra  $\mathcal{A}^*$  obtained in this way is automatically complete in the sense that if  $E \in \mathcal{A}^*$  with  $\mu^*(E) = 0$  and if  $B \subseteq E$ , then  $B \in \mathcal{A}^*$  and  $\mu^*(B) = 0$ . To prove this let A be an arbitrary subset of X and employ Lemma 1.5(3) to observe that  $\mu^*(A) = \mu^*(E) + \mu^*(A) \geq \mu^*(A \cap B) + \mu^*(A \setminus B)$ ; and as before the inequality  $\mu^*(A) \leq \mu^*(A \cap B) + \mu^*(A \setminus B)$  follows from Lemma 1.5(5). Hence B is  $\mu^*$ -measurable and

$0 \leq \mu^*(B) \leq \mu^*(E) = 0$ . We shall now show that in the case that  $\mu$  is a  $\sigma$ -finite measure, it has a unique extension to a measure on  $\mathcal{A}^*$ .

**1.8 Hahn Extension Theorem:**

Suppose that  $\mu$  is a  $\sigma$ -finite measure on an algebra  $\mathcal{A}$ . Then there exist a unique extension of  $\mu$  to a measure on  $\mathcal{A}^*$ .

**Proof:**

The fact that  $\mu^*$  gives a measure on  $\mathcal{A}^*$  proved in theorem 1.7 even without the  $\sigma$ -finiteness assumption. To establish the uniqueness, let  $\nu$  be a measure on  $\mathcal{A}^*$  which agrees with  $\mu$  on  $\mathcal{A}$ .

First suppose that  $\mu$  and therefore  $\mu^*$  and  $\nu$  are finite measures. Let E be any set in  $\mathcal{A}^*$  and let  $(E_n)$  be a sequence

in  $\mathcal{A}$  such that  $E \subseteq \cup E_n$ . Since  $\nu$  be a measure and agrees with  $\mu$  on  $\mathcal{A}$  we have  $\nu(E) \leq \nu\left(\bigcup_{n=1}^{\infty} E_n\right) \leq \sum_{n=1}^{\infty} \nu(E_n) = \sum_{n=1}^{\infty} \mu(E_n)$ . Therefore  $\nu(E) \leq \mu^*(E)$  for any  $E \in \mathcal{A}^*$ . Since  $\mu^*$  and  $\nu$  are additive.  $\mu^*(E) + \mu^*(X \setminus E) = \nu(E) + \nu(X \setminus E)$ . Since the terms on the right hand side are finite and not greater than the corresponding terms on the left hand side, we infer that  $\mu^*(E) = \nu(E)$  for all  $E \in \mathcal{A}^*$ . This establishes the uniqueness when  $\mu$  is a finite measure. Suppose that  $\mu$  is  $\sigma$ -finite and let  $(F_n)$  be an increasing sequence of sets in  $\mathcal{A}$  with  $\mu(F_n) < \infty$  and  $X = \cup F_n$ . Then from the proceeding  $\mu^*(E \cap F_n) = \nu(E \cap F_n)$  for each  $E \in \mathcal{A}^*$ .

Therefore  $\mu^*(E) = \lim \mu^*(E \cap F_n) = \lim \nu(E \cap F_n) = \nu(E)$ , hence  $\mu$  and  $\mu^*$  agrees on  $\mathcal{A}^*$ .

### References:

1. Cohn DL. *Measure Theory*. Birkhauser, Boston 1980.
2. Kingman J, Taylor S. J. *Introduction to Measure and Probability*. Cambridge University Press, 1966.
3. Kirk, R. B. Locally compact ,B-compact spaces, *Indag. Math.* 31, 333–344. Ohta, H., & Tamano, K. (1990), Topological spaces whose Baire measure admits aregular Borel extension, *Trans. Amer. Math. Soc.* 1969; 313:393-415.
4. Wheeler RF. A survey of Baire measures and strict topologies, *Exposition. Math.* 1983, 77 97-190.