



ISSN Print: 2394-7500
ISSN Online: 2394-5869
Impact Factor: 5.2
IJAR 2016; 2(2): 226-228
www.allresearchjournal.com
Received: 18-12-2015
Accepted: 20-01-2016

Dr. Parvati Kushram
Asstt. Prof. of Zoology
Head Deptt. Of Zoology –
Govt. Chandra Vijay College,
Dindori (M.P.).

Physico-chemical studies on Narmada River water at Dindori (M.P.) India

Dr. Parvati Kushram

Abstract

The present paper deals the physicochemical parameters of the river Narmada. Samples were collected season wise from sampling site for analyzing the various physicochemical parameters such as Temperature, pH, TDS, Conductivity, DO, free CO₂, Sulphate, Phosphate, Nitrate, BOD, COD. The work highlights the condition of this river water in various seasons with respect to the parameters mentioned above.

Keywords: Physico-chemical parameters, Narmada River, Dindori.

1. Introduction

In many countries including India, the rivers are not only being exploited but are also used as dumping places for effluents, sewage and solid wastes. Direct or indirect contact of chemicals or waste water to the sources of drinking water cause the undesirable changes in it which becomes dangerous for all living things. Considerable investigations of physico-chemical properties of the river water are carried out in India (Borse, *et al.* 2003^[1], Singh and Gupta, 2004^[2], Barai and Kumar, 2012^[3], Deshmukh, 2012^[4], Chaurasia and Karan, 2013^[5], Kushram, 2013^[6]; Majumder and Dutta, 2014^[7] and Sharma, 2015^[8]).



Photograph showing authoress after collection of water.

A water body affects the environment in its vicinity, like charging of ground water tables, conditions of climate etc. Most of the people like washer man, and fisherman, living in the surrounding area depend on this source of water for their survival. Any damages to this water source by any agency will not only make life miserable but that will also disrupt the aquatic ecosystem. It is therefore necessary to study the quality of river water, on the basis of physico-chemical parameters so as to assess its potability.

2. Material and Methods

A general survey of the river was made for the study of various abiotic parameters. Water has unique property of dissolving and carrying suspension, a huge variety of chemicals, has the undesirable consequence that water can easily become contaminated (APHA, 1989) ^[9]. Water samples were collected monthly in clear glass bottles from surface (max. depth 20 cm)

Correspondence

Dr. Parvati Kushram
Asstt. Prof. of Zoology
Head Deptt. Of Zoology –
Govt. Chandra Vijay College,
Dindori (M.P.)

sites of the river. Water samples were collected in three replicates from surface, column and bottom and mean values of all three observations were taken into consideration. For BOD estimation, water samples were collected separately in dark bottles. The acquisition of meaningful data demands correct sampling and storage procedures. The preservation of samples were done by refrigeration at 4 °C, which is most general accepted method. Water and air temperature were recorded with a digital centigrade thermometer on the date of sampling.

Physico-chemical parameters like water temperature, pH, DO, free Carbon-di-oxide, total alkalinity and conductivity were measured in the field. Other parameters were mostly tested within 24 hrs of collection. A total of 12 limnological parameters of water viz., temperature, turbidity, pH, DO, BOD, COD, Free CO₂, total alkalinity, conductivity, TDS,

Phosphates, nitrate were determined. All the parameters were analysed the standard methods (Golterman, 1969^[10], Michael, 1984^[11], Trivedy and Goel, 1986^[12] and APHA, 1989)^[9] and spectrophotometer SQ 118.


3. Result and Discussion

The present study was conducted at selected sampling station of Narmada River at Dindori town for a period of one year (from Jan 2015 to Dec. 2015). Covering three main seasons i.e. Rainy (July/August/September), Winter (Dec./Jan./Feb.) and Summer (Apr./May/June) in a year.

Physico-chemical and bacteriological parameters were carried out in the samples collected from the study area to study the drinking water quality and pollution level and details of the same was given in table-1 to show the seasonal fluctuations of selected parameters (table-1).

Table 1: Physico-chemical properties of Narmada River for the duration Jan. 2015-Dec.15.

S. No.	Parameters	Rainy Seasons	Winter seasons	Summer seasons	Mean	SD	p
1.	Temp. (°C)	24.8	18.5	28.3	23.87	4.97	16.44
2.	Turbidity	65	95	83	81.00	15.10	152.00
3.	Transparency (cm.)	92	112	87	97.00	13.23	116.67
4.	T.D.S. (mg/l)	378	286	319	327.67	46.61	1448.22
5.	Conductivity	371	277	402	350.00	65.09	2824.67
6.	pH	7.5	8.4	7.6	7.83	0.49	0.16
7.	Alkalinity (mg/l)	134	164	193	163.67	29.50	580.22
8.	D.O. (mg/l)	7.5	9.6	4.3	7.13	2.67	4.75
9.	B.O.D. (mg/l)	12.5	9.8	33.6	18.63	13.03	113.22
10.	C.O.D. (mg/l)	119.5	134.7	70.4	108.20	33.61	752.93
11.	Nitrates (mg/l)	0.023	0.02	0.021	0.02	0.00	0.00
12.	Phosphates (mg/l)	0.045	0.067	0.08	0.06	0.02	0.00
13.	Sulphate	17.2	12.4	17.82	15.81	2.97	5.87



To assess the quality of River, Indian drinking water quality standard IS 10500 (1990) has been adopted. The data harvested during the present study is given in table 1. The present data showed the seasonal variations of all the parameters during the study period.

The water temperature is one of the important parameter in river. In the present study of Narmada River, difference in the fluctuation of water temperature was maximum 28.3 °C to 18.5°C. The season wise studies showed the increased temperature of the river during summer is due to the common effect of intensity of solar radiations, ambient temperature. Decreased values of temperature during winter days are due to low ambient temperature (Gyananath, *et al.* 2000)^[13].

Transparency is light penetration capacity of the water. The color of water is due to concentration of suspended organic

and inorganic particles. Transparency varied from maximum 112 cm to 87 cm. The less transparency observed during rainy season and summer season. While during winter comparatively the water showed more transparency. Low transparency in summer and rainy season in the different water bodies in Jammu (Shashi and Raina, 1990 and Kaushik and Saksena, 1999)^[14, 15].

Total dissolved solids include salt and variety of organic substances, which readily dissolve in water and often impart a degree of hardness. The value of total dissolved solids ranged between max 378 mg/ltr to min 286 mg/ltr. The maximum seasonal value is observed in rainy season and minimum value in winter season. Moreover the low value of total dissolved solids in summer and high in rainy season was also observed (Trivedy, *et al.* 1984)^[16].

The ionic status of water determines the conductivity. During summer and rainy days the enhanced values of conductivity may be due to presence of carbonates and bicarbonates in the water in more quantity due to pollutants. The exchange of inorganic ions by increased micro flora during the winter showed the moderate values of conductivity during the winter period.

Hydrogen ion concentration is considered as an important ecological factor, which is a result of interaction of various substances in water and in numerous biological phenomenon. Nearly neutral pH of water is regulated by carbon dioxide and bicarbonates (Hutchinson, 1957)^[17]. The river water showed well alkaline water through the study period. pH of river ranges between 7.5 to 8.4.

Total alkalinity in river water ranges from 134mg/l to 193mg/l. During the summer season the higher values of hardness were observed and lower values during winter

season (Ugale and Hiware, 1999^[18] and Pratibha, *et al.* 2005^[19]).

Dissolved oxygen is also one of the important factors of water quality, which influences the biota present inside the river water. The seasonal fluctuation of dissolved oxygen in water bodies (Adebisi, 1981)^[20]. Similar pattern of DO observed in the present study, it decreases during the hot days of summer. DO show the inverse relationship with the temperature in river water.

Biological oxygen Demand is a direct measure of O₂ requirement and indirect measure of biodegradable organic matter. The maximum B.O.D. was 33.6mg/l in summer and minimum 9.8mg/l in winter (Seenaya and Zafar, 1979)^[21].

Chemical Oxygen Demand indicates the extent of chemical pollution mainly from industrial effluents. The C.O.D. values observed maximum in winter and minimum in rainy season (Singh and Roy, 1995)^[22]. In the present study, nitrate values ranged between 0.02 to 0.023 mg/l maximum of nitrate values were reported during rainy season and summer and minimum during winter. The high value of nitrate in rainy season, which is linked to heavy run-off of the organic matter from the catchments (Gohram, 1961)^[23].

Phosphate concentration in river water ranged between 0.045 mg/l to 0.08 mg/l. more concentration of phosphates recorded during the summer and winter season. It may be due to deposition of ashes and bones under religious activities and decomposition of organic matter in the water sediments.

Sulphate is produced by biological oxidation of sulphur content of organic matter. The sulphate value ranges between 12.4 mg/l to 17.82 mg/l. Minimum concentration of sulphate was in winter and maximum in summer (Angadi, *et al.* 2005)^[24].

4. Conclusion

From the above study, it may conclude that except little variation, all the physico-chemical parameters were in permissible limit at the study site of the Narmada River. It is suggested that proper measures are necessary to avoid contamination as water is used for drinking purpose. At present the river is suitable for irrigation and fishery purpose.

5. Acknowledgement

I am thankful to the authority of college for granting permission to carry out this work.

6. References

- Borse SK, Lohar PS, Bhave PV. Hydrobiological study of algae of Aner River, Jalgaon (Maharashtra), *J Aqu Bio.* 2003; 18(1):15-18.
- Singh M, Gupta KC. Physico-chemical studies of water of river Yamuna at Mathura, *Ecol. Envi. And Cons.*, 2004; 10(2):193-196.
- Barai SR, Kumar Satish. Evaluation of the physicochemical characteristics of river Varuna at Varanasi, India, *J Environ Biol.* 2012; 34:259-265.
- Deshmukh BS. Hydrobiological study of river Pravara in Ahmednagar district (Maharashtra), *Bionano Frontier, Eco Revolution 2012 Colombo*, 2012, 89-92.
- Chaurasia S, Karan Raj. Water quality and pollution load of river Mandakini at Chitrakoot, India, *Int. Res. J Environ Sci.* 2013; 2(6):13-19.
- Kushram Parvati. Study of physico-chemical and biological analysis of drinking water of Dindori town with special reference to quality status Unpublished thesis A.P.S. University, Rewa (M.P.), 2013.
- Majumder S, Dutta TK. Studies on seasonal variations in physico-chemical parameters in Bankura segment of the Dwarakeshwar River (W.B) India, *IJAR*, 2014; 2(3):877-881.
- Sharma Pankaja. Seasonal Variations in Physico-Chemical Properties of Narmada River in Dindori Madhya Pradesh, India, *International Journal for Research in Applied Science & Engineering Technology (IJRASET)*, 2015; 3(XII):285-288.
- APHA. Standard methods for the examination of water and wastewater. 17th edition, American Public Health Association, Washington D.C, 1989.
- Golterman HL. (Ed). *Methods of chemical analysis of freshwaters.* IBP Handbook Blackwell, Oxford, 1969, 8.
- Michael P. *Ecological methods for field and laboratory investigations.* Tata-McGraw Hill Pub. Com. Ltd., New Delhi, 1984, 404.
- Trivedy PK, Goel RK. *Chemical and Biological methods water pollution studies,* Environmental publication Karad India, 1986.
- Gyananath G, Shevnikar SV, Samiuddin S. Water quality analysis of river Godavari during Holimela at Nanded, *Poll. Res.*, 2000; 19(4):673.
- Shashi Kant, Raina AK. Limnological studies of two ponds in Jammu. *J Env Biol.* 1990; 11(2):137-144.
- Kaushik S, Saksena DN. Physico-chemical limnology of certain water bodies of central India. In: *fresh water ecosystems of India* (Ed. K. Vijay Kumar) Daya Publ., House, Delhi., 1999, 1-58.
- Trivedy RK, Goel PK, Shrotri AC, Khatavkar SD. *Prospective in Aquaculture, Biol.*, 1984, 15-18.
- Hutchinson GE. *A treatise on limnology* John Wiley and Sons, New York, 1957; II:1015.
- Ugale BJ, Hiware CJ. Limnological study of an ancient reservoir Jagtunga Samudra located at Kandhar, Dist. Nanded, Maharashtra, India. *Eco. Envi and Cons.*, 1999; 11(3-4):473-475.
- Pratibha Raithak V, Bhuktar AS. Physicochemical analysis of drinking water from different areas of Aurangabad, *Poll. Res.*, 2005; 24(3):727-728.
- Adebisi. The physicochemical hydrobiology of a tropical river upper Ogun river Nigeria. *Hydrobiol*, 1981; 79(2):157-165.
- Seenaya G, Zafar AR. An ecological study of Mir Alam Lake, Hyderabad, India. *Indian J Biol.* 1979; 11(2):330-335.
- Singh JP, Roy SP. Limnobiological investigation of Karwarlake, Begusarai, Bihar *Env. Eco.*, 1995; 13:330-335.
- Gohram. The chemical composition of some waters from Dune slacks at Sandscale, North Lancshire, *J Ecology.* 1961; 49(1):79-82.
- Angadi SB, Shiddamallayya N, Patil PC. Limnological studies of Papanash pond, Bidar, Karnataka, *J Env Biol.* 2005; 26(2):213-216.