



ISSN Print: 2394-7500
ISSN Online: 2394-5869
Impact Factor: 5.2
IJAR 2016; 2(8): 323-325
www.allresearchjournal.com
Received: 20-06-2016
Accepted: 21-07-2016

Katari Kantha
Department of Community
Health Nursing, Narayana
College of Nursing, Nellore,
Andhra Pradesh, 524003,
India.

Arumugam Indira
Principal, Narayana College of
Nursing, Nellore, Andhra
Pradesh, 524003, India.

Prevalence of hypertension among the adults in Mahalakshmpuram vs Allipuram

Katari Kantha and Arumugam Indira

Abstract

Background: High blood pressure (BP) is ranked as the third most important risk factor for attributable burden of disease in south Asia (2010). Hypertension (HTN) exerts a substantial public health burden on cardiovascular health status and healthcare systems in India.

Aim: to assess the prevalence of hypertension

Setting and Design: The study was conducted in Mahalakshmpuram (coastal area) and Allipuram (non-coastal area) by using a descriptive design.

Materials and Methods: A total of 500 samples were included in this study. Among this, 250 samples belongs to coastal area and 250 samples belongs to non-coastal area by using convenience sampling technique.

Statistical Analysis Used: The collected data was organized, tabulated, analyzed and interpreted by using descriptive and inferential statistics based on the objectives of the study.

Results: In Mahalakshmpuram, Out of 250 samples, With regard to the category of the blood pressure 16(6.4%) had stage-I hypertension, 2(0.8%) had stage-II hypertension, 0(0.00%) had stage-III hypertension, 39(15.6%) had grade-I isolated systolic hypertension, and 2(0.8%) had grade-II isolated systolic hypertension. Known Hypertensive cases are 39(15.6%), Newly diagnosed cases are 20(8.00%). With regard to BMI, among 250 samples 29(11.6%) were overweight and 8(3.2%) were obese. in Allipuram, among 250 samples, 141(56.4%) had stage-I hypertension, 18(7.2%) had stage-II hypertension, 0(0.00%) had stage-III hypertension, 54(21.6%) had grade-I hypertension, and 14(5.6%) had grade-II hypertension. Known Hypertensive cases are 151(60.4%). Newly diagnosed cases are 47(18.8%). With regard to BMI among 250 samples 43(17.2%) were overweight and 37(14.8%) were obese.

Conclusion: The above results shown that blood pressure values are high in the Allipuram (non-coastal area) than Mahalakshmpuram (coastal area).

Keywords: Hypertension, non-coastal area, coastal area, heart attack, stroke

Introduction

One in three adults worldwide has high blood pressure. Hypertension increases the risk of heart attack, stroke, kidney failure and much other associated co morbidity. Treating raised blood pressure and maintaining it below 140/90 mmHg is associated with a reduction in cardiovascular complication. The theme for World Health Day (WHD) 2013 is "high blood pressure". The goal of WHD 2013 is to reduce heart attacks and strokes. Keeping in line with the WHO, Government of India, Country Cooperation Strategy, the WHO 2013 events in India are aimed at raising the awareness amongst national policymakers, program managers and other stakeholders on the need to strengthen the Indian health system to make it competent enough to respond to hypertension and related co morbidities [1].

Kantha, K and Indira, A. (2015) conducted a cross sectional study on prevalence of hypertension among the adults in coastal and non-coastal areas. A total of 5000 samples were included in the study. In that 2500 samples belongs to coastal areas and 2500 samples belongs to non-coastal areas. The prevalence of stage-I hypertension in coastal areas is 460(18.4%) but in non-coastal areas it is 1413(56.50%). The results indicate that there is high prevalence of hypertension in non-coastal areas than coastal areas [2].

Correspondence
Katari Kantha
Department of Community
Health Nursing, Narayana
College of Nursing, Nellore,
Andhra Pradesh, 524003,
India.

Arumugam Indira *et al.* (2015) conducted a study on prevalence of prehypertension among the adults in coastal and non-coastal areas. The study results shown that regarding prehypertension in SBP, in coastal areas 1129(45.16%) and in non-coastal areas 971(38.84%). The results indicate that there is high prevalence of pre hypertension in coastal areas than non-coastal areas. Further studies are needed to find out the reasons and measures to control high blood pressure is necessary [3].

Even today there is scarcity of the studies in coastal and non-coastal areas of India. With this background, present study has been undertaken to study the prevalence of hypertension.

Objectives of the Study

- To assess the prevalence of hypertension among adults of coastal and non-coastal areas.
- To identify the risk factors of hypertension among adults of coastal and non-coastal areas.
- To compare the prevalence of hypertension between coastal and non-coastal areas.
- To find association between the prevalence of hypertension with selected socio demographic variables.

Detailed Research Plan

Research Approach: Quantitative Approach.

Research Design: Descriptive design.

Research Setting: The study was conducted in Mahalakshmiapuram (coastal area) and Allipuram (non-coastal area) by using a descriptive design.

Coastal area means areas within 2km from mean low water mark (MLWM) or mean high water mark (MHWM).

Non-coastal area means areas far 2km from mean low water mark (MLWM) or mean high water mark (MHWM).

Sampling Technique: Convenience sampling technique

Sample Size: A total of 500 samples were included in this study. Among this, 250 samples belongs to Mahalakshmiapuram (coastal area) and 250 samples belongs to Allipuram (non-coastal area).

Results and discussion

Comparison of Blood Pressure in Mahalakshmiapuram and Allipuram

Table 1: Comparison of Blood Pressure in Mahalakshmiapuram and Allipuram.

Blood Pressure Category	Mahalakshmiapuram		Allipuram		Correlation coefficient	Standard deviation
	(f)	(%)	(f)	(%)		
Optimal	34	13.6%	0	0.00%	0.06	40.01
Norma	62	24.8%	4	1.6%		
High Normal	95	38%	48	19.2%		
Stage-I	16	6.4%	141	56.4%		
Stage-II	2	0.8%	18	7.2%		
Stage-III	0	0.00%	0	0.00%		
Grade-I	39	15.6%	36	14.4%		
Grade-II	2	0.8%	3	1.2%		




Fig 1: Comparison of Blood Pressure in Mahalakshmiapuram and Allipuram

The prevalence of stage-I BP in coastal area is 16(6.4%) but in non-coastal areas it is 141(56.4%). The correlation coefficient value is highly significant (0.06) and the standard deviation is 40.01.

Comparison of Body Mass Index in Mahalakshmiipuram and Allipuram




Fig 2: Comparison of Body Mass Index in Mahalakshmiipuram and Allipuram

The prevalence of overweight samples in coastal area is 29 (11.6%), obesity is 8(3.2%) but in non-coastal areas it is 43(17.2%) and 37(14.8%). The correlation coefficient value is highly significant (0.99) and the standard deviation is 81.46.

Association of Socio Demographic Data with the Blood Pressure in Mahalakshmiipuram

There is a significant association of demographic variables with Age, Amount of oil used for per day, intake of fish, and are you a known hypertensive and remaining are non-significant.

Association of Socio Demographic Data with the Blood Pressure in Allipuram

There is a significant association of demographic variables with Age, Family, Type of house, sleeping Hours, Food pattern, Amount of oil used for per day, Amount of vegetables per day, Habits and Entertainment and remaining are non-significant.

Conclusion

- The above results shown that grade-1 and grade-2 isolated systolic hypertension values are higher in the Allipuram (non-coastal area) than in the Mylapuram (coastal area).
- Among hypertension cases the prevalence of obesity is more in non-coastal area than coastal area.
- The variables like Age, exercise, Type of oil used for cooking, Type of salt used, habits, intake of fish, are you having stress and are you a known hypertensive are the influencing risk factors for the development of hypertension among the adults.

References

1. Anchala R, Kannuri NK, Pant H. Hypertension in India: a systematic review and meta-analysis of prevalence, awareness, and control of hypertension. *J Hypertens*. 2014; 32(6):1170-7.
2. Katari Kantha, Arumugam Indira. Prevalence of hypertension among the adults in coastal and non-coastal areas *International Journal of Development Research*. 2015; 05(01):3134-3139,
3. Arumugam Indira. Prevalence of Pre Hypertension among the Adults Aged 20-60 Years in Coastal and Non Coastal areas *International Journal of recent scientific research*. 2015; 6(11):7166-7170,
4. Arumugam Indira, Katari Kantha. Nutritional status of antenatal and postnatal mothers in selected rural areas at Nellore. *International Journal of Recent Scientific Research*. 2015; 6(2):2796-2800,
5. Katari Kantha. Knowledge of the Expectant Fathers regarding safe Motherhood in Narayana Medical College Hospital, Nellore. *International Journal of Recent Scientific Research*. 2015; 6(4):3423-3427.
6. Vanaja Kumari B. Comparative Study to Assess the Knowledge Regarding Behavioural Problems of Underfive Children Among Employed and Unemployed Mothers. *International Journal of Recent Scientific Research*. 2015; 6(5):4183-4187.
7. Katari Kantha. Nutritional Status of Infant Children in Selected Rural Areas at Nellore *International Journal of Recent Scientific Research*. 2015; 6(6):4597-4601