



ISSN Print: 2394-7500
 ISSN Online: 2394-5869
 Impact Factor: 5.2
 IJAR 2016; 2(9): 719-720
 www.allresearchjournal.com
 Received: 10-07-2016
 Accepted: 11-08-2016

A Vijayasankar
 Assistant Professor,
 Department of Mathematics,
 National College, Trichy-
 620001, Tamil Nadu, India

MA Gopalan
 Professor, Department of
 Mathematics, Shrimati Indira
 Gandhi College, Trichy-620002,
 Tamil Nadu, India

V Krithika
 Research Scholar, Department
 of Mathematics, National
 College, Trichy-620001, Tamil
 Nadu, India

Correspondence
A Vijayasankar
 Assistant Professor,
 Department of Mathematics,
 National College, Trichy-
 620001, Tamil Nadu, India

Special Dio-quadruples with property D(2)

A Vijayasankar, MA Gopalan and V Krithika

Abstract

This paper concerns with the study of constructing a special Dio quadruples (a,b,c,d) such that the product of any two elements of the set minus with their sum and increased by two is a perfect square.

Keywords: Diophantine quadruples, pell equation. 2010 mathematics subject classification 11D99.

Introduction

The problem of constructing the sets with property that the product of any two of its distinct elements is one less than a square has a very long history and such sets were studied by Diophantus [1]. A set of m positive integers $\{a_1, a_2, \dots, a_m\}$ is said to have the property D(n), $n \in \mathbb{Z} - \{0\}$ if $a_i a_j + n$, a perfect square for all $1 \leq i < j \leq m$ and such a set is called a Diophantine m-tuples with property D(n). Many mathematicians considered the construction of different formulations of Diophantine quadruples with the property D(n) for any arbitrary integer n and also for any linear polynomials in n. In this context, one may refer [2-13] for an extensive review of various problems on Diophantine quadruples. This paper aims at constructing special dio-quadruples where the special mention is provided because it differs from the earlier one and the special Dio-quadruples is constructed where the product of any two members of the quadruples with the subtraction of the sum of the same members and the addition of two satisfies the required property.

Method of Analysis

Special Dio-Quadruple with Property D(2)

Let $a(n) = 3^n$ and $b(n) = 3^n + 2$ be two integers such that $ab - (a + b) + 2$ is a perfect square.

Let $C_N(n)$ be any non-zero integer such that

$$(3^n - 1)C_N(n) - 3^n + 2 = p_N^2(n) \tag{1}$$

$$(3^n + 1)C_N(n) - (3^n + 2) + 2 = q_N^2(n) \tag{2}$$

Eliminating $C_N(n)$ from (1) and (2), we obtain

$$(3^n + 1)p_N^2(n) - (3^n - 1)q_N^2(n) = 2 \tag{3}$$

$$\text{Setting } p_N(n) = X_N(n) + (3^n - 1)T_N(n) \tag{4}$$

$$q_N(n) = X_N(n) + (3^n + 1)T_N(n) \tag{5}$$

in (3), we get

$$X_N^2(n) = (3^{2n} - 1)T_N^2(n) + 1 \tag{6}$$

whose general solution is

$$X_N(n) = \frac{1}{2} \left[(3^n + \sqrt{3^{2n} - 1})^{N+1} + (3^n - \sqrt{3^{2n} - 1})^{N+1} \right] \tag{7}$$

$$T_N(n) = \frac{1}{2\sqrt{3^{2n} - 1}} \left[(3^n + \sqrt{3^{2n} - 1})^{N+1} - (3^n - \sqrt{3^{2n} - 1})^{N+1} \right]$$

Substituting $N=0$ in (7) and (4) and employing (1), we get,

$$C_0(n) = 1 + 4 * 3^n$$

Note that the triple $(a, b, C_0(n))$ is a special Dio-triple with property D(2)

Again, substituting $N=1$ in (7) and (4) and employing (1), we get,

$$C_1(n) = 16 * 3^{3n} - 4 * 3^n + 1$$

It is seen that the $(a, b, C_0(n), C_1(n))$ is a special Dio-Quadruple with property D(2).

Again, substituting $N=2$ in (7) and (4) and employing (1) we get,

$$C_2(n) = 1 + (8 * 3^{3n} - 4 * 3^{2n} - 4 * 3^n + 2)(8 * 3^{2n} + 4 * 3^n)$$

It is seen that the $(a, b, C_1(n), C_2(n))$ is a special Dio-Quadruple with property D(2).

And also, substituting $N=3$ in (7) and (4) and employing (1) we obtain,

$$C_3(n) = 1 + (256 * 3^{7n} - 320 * 3^{5n} + 112 * 3^{3n} - 8 * 3^n)$$

It is seen that the $(a, b, C_2(n), C_3(n))$ is a special Dio-Quadruple with property D(2). The repetition of the above process leads to the result that the quadruple $(a, b, C_{N-1}(n), C_N(n))$.

$N=1, 2, 3, 4, \dots$ is a special Dio-Quadruple with property D(2).

Some numerical examples are tabulated below:

Table. Special Dio-Quadruple With Property D(2)

n	(a, b, C_0, C_1)	(a, b, C_1, C_2)	(a, b, C_2, C_3)
1	(3,5,13,421)	(3,5,421,14281)	(3,5,14281,485113)
2	(9,11,37,11629)	(9,11,11629,3744217)	(9,11,3744217,1205625961)
3	(27,29,109,314821)	(27,29,314821,917385481)	(27,29,917385481,2673260974000)
4	(81,83,325,8502733)	(81,83,8502733,223128693100)	(81,83,223128693100,5855343157000000)
5	(243,245,973,229581541)	(243,245,229581541,54225782260000)	(243,245,54225782260000,12807804410000000000)

5. Conclusion

This paper concerns with the construction of special dio - quadruples with property D(2). One may search for special dio-quadruples consisting of special numbers with suitable property.

6. References

1. Abu Muriefah FS, Al-Rashed A. Some Diophantine quadruples in the ring $Z[\sqrt{-2}]$ Math. Commun, 2004; 9:1-8.
2. Assaf E, Gueron S. Characterization of regular Diophantine quadruples, Elem. Math. 2001; 56:71-81.
3. Bashmakova IG. (ed.), Diophantus of Alexandria, Arithmetic's and the book of polynomial numbers, Nauka Moscow, 1974.
4. Dujella A. Some polynomial formulas for Diophantine quadruples, Grazer Math. Ber. 1996; 328:25-30.
5. Dujella A. Some estimates of the number of Diophantine quadruples, Publ. Math. Debrecen, 1998; 53:177-189.
6. Filipin A, Fujita Y. Any polynomial D (4)-quadruple is regular, Math. Commun, 2008; 13:45-55.
7. Filipin A. An irregular D (4)-quadruple cannot be extended to a quintuple, Acta Arith, 2009; 136:167-176.
8. Franusic Z. Diophantine quadruples in the ring $Z[\sqrt{-2}]$ Math. Commun, 2004; 9:141-148.
9. Franusic Z. Diophantine quadruples in Quadratic Fields, Dissertation, University of Zagreb, 2005.
10. Gopalan MA, Srividhya G. Diophantine quadruples for Fibonacci and Lucas numbers with property D (4). Diophantus J Math. 2012; 1:15-18.
11. Gibbs P. Diophantine quadruples and Cayley's hyperdeterminant, XXX Mathematics Archive math. NT/0107203
12. Gibbs P. Adjugates of Diophantine Quadruples Integers. 2010; 10:201-209.
13. Srividhya G. Diophantine quadruples for Fibonacci numbers with property D (1). Indian journal of

Mathematics and Mathematical Sciences. 2009; 5:57-59.