



ISSN Print: 2394-7500  
 ISSN Online: 2394-5869  
 Impact Factor: 5.2  
 IJAR 2017; 3(1): 963-967  
 www.allresearchjournal.com  
 Received: 16-11-2016  
 Accepted: 20-12-2016

**Sateydeo Singh**  
 Research Scholar,  
 Department of Chemistry  
 S.K.M.U, University Dumka,  
 Jharkhand, India

## Study of reduction products of ortho-hydroxyl substituted azo dyes on biodecolorization of azo dyes

**Sateydeo Singh**

### Abstract

In this paper, the mediated effects of reduction products of some ortho-hydroxyl substituted azo dyes on biodecolorization were investigated. The results indicated that the addition of reduction products could effectively accelerate dye decolorization by *Shigella* sp. QRZ-1. The best accelerating effect was obtained with the addition of reduction products of Acid Red 14 (AR14), resulting in an over 3-fold increase in decolorization efficiency of many azo dyes. In sequencing batch reactor experiments, the accelerating effect of reduction products of AR14 was more obvious (1.5-fold) during the startup of the system. When the dye concentration was increased to 500 mg L<sup>-1</sup>, the accelerated decolorization efficiency was still maintained around 95%. The presence of AR14 in the feed enhanced the decolorization performance of anaerobic sludge, indicating that the strategy may be beneficial for practical application. 1-Naphthol-2-amino-4-sulfonic acid, which is one of the reduction products of AR14, may function as redox mediator to speed up azo dye biodecolorization.

**Keywords:** ortho-hydroxyl, azo dyes and biodecolorization

### Introduction

The mechanism of azo dye reduction is still a subject of debate. Some suggested that it should be a specific reaction by intracellular azoreductase while others reported it to be an extracellularly unspecific reduction process. Intracellular azoreductase cannot reduce all types of azo dyes, especially those sulfonated azo dyes with high polarity and polymeric azo dyes with high molecular weight, which have limited membrane permeability.

The extracellular reduction of the azo dye is usually a slow process. However, it could be significantly enhanced by the use of redox mediators such as lawsone, riboflavin, and anthraquinone-2,6-disulfonate (AQDS) [1-3]. Quinone compounds used as redox mediators are reduced by cytoplasmic or membrane-bound quinone reductase. The hydroquinones formed can then reduce azo dyes in a purely chemical redox reaction [4]. However, the continuous dosing of redox mediators implies continuous expenses related to procurement of the chemical, as well as continuous discharge of biologically recalcitrant compounds resulting in secondary contamination. In addition, it is usually difficult to immobilize quinone redox mediators. The catalytic effects of immobilized mediators were not satisfying [5].


It was reported that monoazo dye Acid Orange 7 (AO7) has an autocatalysis effect. 1-Amino-2-naphthol, the constituent aromatic amine and degradation product of AO7 can accelerate the reduction process, possibly by mediating the transfer of reducing equivalents [6, 7]. Similarly, it was reported that the decolorized supernatants of Direct Blue 71 and Direct Blue 53, two tri-azo dyes, increased the biodecolorization rate of Methyl Orange. Such mediators can be generated during azo dye reduction, and mineralized by sequential aerobic treatment without secondary contamination. However, the rate constant of AO7 reduction mediated with 1-amino-2-naphthol was 10-fold lower than that obtained using AQDS. Thus some better mediators generated from azo dye reduction are expected for practical application.

In this study of reduction products of some ortho-hydroxyl substituted azo dyes on biodecolorization of azo dyes were investigated.

### Materials and methods

**Chemicals:** Azo dyes used in this study are shown in Fig. 1.

**Corresponding Author:**  
**Sateydeo Singh**  
 Research Scholar,  
 Department of Chemistry  
 S.K.M.U, University Dumka,  
 Jharkhand, India



**Fig 1:** Azo dyes used in this study

All the dyes were purchased from Tianjin Tianshun Chemical Dyestuff Co., ed in this study. Ltd. Other chemicals used in the study were all purchased from Sigma.

**Preparation of the reduction products of ortho-hydroxyl substituted azo dyes:** Cell suspension (5 mL) was inoculated to 135 mL serum bottles filled with sterilized decolorization medium containing 1 M specific acid azo dye. Then the vials were sealed with butyl rubber stoppers, and statically cultivated at 30°C until the complete decolorization of dyes. Cells were removed by centrifugation (24,000 x g, 10 min), and the supernatants were kept at 4°C before used as mediator in the following decolorization studies. The exact amounts of specific

mediators in the supernatants were determined with high performance liquid chromatography before use.

**Batch decolorization experiments:** All batch decolorization experiments were conducted in 135 mL serum bottles filled with decolorization medium. Condensed cell suspensions or mixed sludge were transferred under anaerobic conditions to the bottles to reach  $OD_{560} = 0.4$ , or sludge concentration (MLSS) of  $3.0 \pm 0.1 \text{ g L}^{-1}$ . Azo dyes and reduction products used as mediators were added by syringe, respectively. The serum bottles were sealed with butyl rubber stoppers, and then statically cultivated at 30 °C. All the experiments were done in triplicate.

**Analytical methods:** Absorbance was measured spectrophotometrically with spectrophotometer Uasco V-560) at respective maximum wavelength of azo dyes. The concentration was calculated according to the corresponding calibration curves.


The cyclic voltammetric experiments were carried out using a PARSTAT 2273 electrochemical system, employing a three-electrode configuration: glassy carbon electrode (d = 3 mm) as the working electrode, a platinum wire as the auxiliary electrode, and saturated calomel electrode as the reference electrode. As electrolyte support, 0.1 M of Na<sub>2</sub>SO<sub>4</sub> solution was used. The cyclic voltammetric scan was conducted at a rate of 100 mV s<sup>-1</sup> within the potential interval from -1200 to 1200 mV. The test solutions were flushed with high-purity N<sub>2</sub> for 30 min to remove interfering oxygen.

The aromatic amines were detected with high performance liquid chromatography/mass spectrometry (HPLC/MS, Shimadzu HPLC-2010A system). The chromatographic column is Shim-pack VP-ODS (150 mm x 2.0 mm, C18 column). The column temperature was 40°C. For 1-naphthylamine-4-sulfonic acid, the mobile phase consisted of methanol and water containing dibutylamine (2.5 mM) and acetic acid (2.5 mM). The elution program began with 30% methanol and 70% water. The percentage of methanol was linearly increased to 70% over 10 min. The flow rate was 0.2 mL min<sup>-1</sup>. For 1-naphthol-2-amino-4-sulfonic acid,

the mobile phase consisted of methanol and water containing dibutylamine (4 mM). The elution program began with 60% methanol and 40% water. The percentage of methanol was linearly increased to 100% over 40 min. The flow rate was 1.0 mL min<sup>-1</sup>. MS was performed under the following conditions: electrospray ionization source (ESI-), detector pressure 1.6 V, scan mode.

## Results and discussion

**Characteristics of strain QRZ-1:** A pure quinone-reducing isolate QRZ-1 was obtained from the sludge mixture after 2 months of enrichment at 30 °C in the growth medium. It is a Gram-negative, non-motile, rod-shaped, anaerobic bacterium (0.2-0.3 μm wide and 0.7-0.9 μm long). Colonies are mucoid, rounded and white when cultivated on solid growth medium. It is catalase-positive and produces acids from glucose. It can use glucose, maltose, sucrose, galactose, fructose, mannose, oxalate, lactate, n-xylose, α-lactose and sorbose as sole carbon source. However, citrate, acetate, malonate and succinate cannot be utilized by the strain. It is resistant to tetracycline and streptomycin. On the basis of 16S rDNA sequences comparison at NCBI and RDP II, QRZ-1 has higher homology (99%) with the genus *Shigella* within the family Enterobacteriaceae of the gamma subclass of the Proteobacteria. The phylogenetic analysis demonstrated that QRZ-1 is distinct from other *Shigella* species (as shown in Fig. 2).




**Fig 2:** Phylogenetic tree of strain QRZ-1 based on the 16S rRNA sequences using the MP method. The sources of sequences are shown as GenBank accession number in brackets. Only bootstrap (500 replicates) values above 50% are shown

According to the physiological and biochemical characteristics and 16S rRNA gene sequencing results, the strain QRZ-1 is a close relative of the genus *Shigella* and designated as *Shigella* sp. strain QRZ-1.

It has been found that many microorganisms are capable of reducing quinones, such as Fe(III)-reducers *Pantoea agglomerans* SP1, *Geobacter sulfurreducens*, and *Shewanella decolorationis* S12, uranium-reducer *Deinococcus radiodurans*, sulfate-reducer *Desulfovibrio G11*, halo-respiring bacterium *Desulfobacterium dehalogenans*, fermentative bacterium *Propionibacterium freudenreichii*, and methanogenic archaea *Methanospirillum hungatei* [8-11]. However, to our knowledge, this is the first report on quinone reduction by genus *Shigella*. Genes encoding putative quinone

oxidoreductases have been reported in genome of *Shigella* genus [12]. Previous studies using other bacterial strains demonstrated that quinoid redox mediators can be reduced by cytoplasmic or membrane-bound quinone reductase to speed up azo dye decolorization. Thus QRZ-1 was used in the following mediated decolorization studies.


**Decolorization experiments with strain QRZ-1:** The effects of the reduction products of A07, AR73, AR18 and AR14 (named RA07, RAR73, RAR18 and RAR14, respectively) on decolorization of RR120 by strain QRZ-1 were studied. During the decolorization process we investigated, QRZ-1 kept its normal growth and almost no obvious growth difference was observed in the presence of different mediators. As shown in Fig. 3,



**Fig 3:** Growth of strain QRZ-1 ( $OD_{660}$ ) and biodecolorization of RR120 by strain QRZ-1 in the presence of AQDS1 (0.06 mM), AQDS2 (0.12mM) and 0.12mM reduction products of RAR14, RAO7, RAR18 and RAR73

without the addition of reduction products, only 17% RR120 was removed in 10 h. On the other hand, the addition of reduction products of acid azo dyes accelerated the decolorization of RR120 effectively. The addition of RAR14 resulted in removal of 91% RR120 in 10 h, which was the best accelerating effect and better than that obtained with the model redox mediator AQDS.

Two current peaks of reversible oxidation and reduction were observed at potentials of approximately 338 and -209 mV, respectively which indicates the reversible redox process of RAR14. In contrast, no peak was found in voltammograms measured when no reduction products were formed at 0 min (As shown in Fig. 4).




**Fig 4:** Cyclic voltammograms of the system at 0 min (blue line) and when RAR14 was formed (red line). (For interpretation of the references to color in this figure legend, the reader is referred to the web version of the article.)

Thus RAR14 could function as redox mediator and was responsible for the accelerated decolorization performance. Similar current peaks were also observed with other reduction products (data not shown). It was reported that two flavoproteins from *S. dysenteriae* could act as azoreductase for the decolorization of some azo dyes in vitro [13]. However, due to their high polarity and large molecular weight, azo dyes such as RR120 are unlikely to

pass through the cell membrane. The decolorization of RR120 might be a process occurred extracellularly. The reduction products of these ortho-hydroxyl substituted azo dyes possibly act like AQDS and RAO7 as redox mediators by mediating the transfer of reducing equivalents from the cell to extracellular azo dyes.

Effects of RAR14 concentration on decolorization of RR120 by strain QRZ-1 were investigated (As shown in Fig. 5).




**Fig 5:** Effects of RAR14 concentration (0-0.4 mM) on biodecolorization of RR120

The results indicated that the decolorization efficiency was enhanced dramatically with the increase of RAR14 concentration. A 3.5-fold increase of decolorization efficiency was obtained with the addition of 0.05 mM RAR14. The addition of 0.1-0.2 mM RAR14 resulted in approximately 4.5-fold increase. However, with further increase of the RAR14 concentration over 0.3 mM, the stimulating effect on decolorization decreased correspondingly. When AQDS is used as redox mediator for azo dye decolorization, the enzymatic reduction of the AQDS is generally suggested to be the rate-limiting step. Similar decrease of accelerating effects due to higher concentration of AQDS were reported previously [14].


Since the thermodynamics of different electron-donating half-reactions varied, the reaction rate may be influenced by the type of electron donor. Different carbon sources (0.5 g L<sup>-1</sup>) were therefore chosen to determine their effects on RR120 decolorization by strain QRZ-1 in the presence of RAR14 (0.12 mM). The strain was able to grow on all tested carbon sources, whereas the decolorization activities varied. RR120 decolorization was observed after 10-h cultivation with glucose, sucrose, maltose, mannose, oxalate, lactate, and peptone as carbon sources, respectively. The maximum decolorization efficiency (91.3%) was obtained with glucose. Using lactate and oxalate as carbon sources resulted in less than 10% decolorization efficiency.

It was shown that AQDS can be used as redox mediator to accelerate the decolorization of many azo dyes. Most of the azo dyes investigated in this study are characteristic of complex structure and high molecular weight (Fig. 1). It is believed that they cannot enter into the bacterial cells and are difficult to be biodecolorized. As shown in Fig. 6, the addition of RAR14 could dramatically enhance the decolorization of these different azo dyes by strain QRZ-1.



**Fig 6:** Comparison of accelerating effects of RAR14 (0.12 mM) and AQDS (0.12 mM) on decolorization of different azo dyes by strain QRZ-1. Cultures without the addition of redox mediator served as control

**Decolorization experiments with anaerobic sludge:** SBR experiments with anaerobic sludge also demonstrated the enhancement of RR120 decolorization in the presence of RAR14 (As shown in Fig. 7).



**Fig 7:** Effects of RAR14 on RR120 decolorization by anaerobic sludge in SBR

The accelerating effect was more obvious (1.5-fold higher) during the startup of the system. When RR120 concentration was increased to 500 mg L<sup>-1</sup>, the decolorization efficiency was still maintained at approximately 95%, and the abiotic adsorption of dyes on the biomass was insignificant (data not shown). Moreover, the decolorization of mixed dyes by anaerobic sludge was also speeded up in the presence of AR14 (data not shown). Thus RAR14 may act as redox mediator for various microorganisms in the sludge and be beneficial for the practical bioprocess of wastewater treatment.

### Conclusions

This study demonstrated that the reduction products of some ortho-hydroxyl substituted acid azo dyes, such as 1-naphthol-2-amino-4-sulfonic add formed in AR14 decolorization by *Shigella* sp. strain QRZ-1 or anaerobic sludge could function as redox mediator and accelerate azo dyes biodecolorization. Compared to the synthetic mediator AQDS, RAR14 demonstrated better mediating effects. In addition, estimated from the results obtained in our studies,

the use of metabolites generated from biological reduction possesses will save around \$120-160 per ton of dye wastewater. The addition of such ortho-hydroxyl substituted acid azo dyes in the feed may be a promising and cost-effective strategy to improve the biodecolorization of azo dye wastewater.

### References

1. Van der Zee FP, Bouwman RH, Strilk DP, Lettinga G, Field JA. Application of redox mediators to accelerate the transformation of reactive azo dyes in anaerobic bioreactors, *Biotechnol Bioeng* 2001;75:691-701.
2. Rau J, Knackmuss HJ, Stolz A. Effects of different quinoid redox mediators on the anaerobic reduction of azo dyes by bacteria, *Environ Sci Technol* 2002;36:1497-1504.
3. Encinas-Yocupicio AA, Raw-Flores E, Sanchez-Diaz F, Dos Santos AB, Field JA, Cervantes FJ. Catalytic effects of different redox mediators on the reductive decolorization of azo dyes, *Water Sci Technol* 2006;54:165-170.
4. Kuddich M, Keck A, Klein J, Stolz A. Localization of the enzyme system involved in anaerobic reduction of azo dyes by *Sphingomonas* sp. strain BN6 and effect of artificial redox mediators on the rate of azo dye reduction, *Appl. Environ. Microbiol* 1997;63:3448-3455.
5. Guo J, Zhou J, Wang D, Tian C, Wang P, Uddin MS, Yu H. Biocatalyst effects of immobilized anthraquinone on the anaerobic reduction of azo dyes by the salt-tolerant bacteria, *Water Res* 2007;41:426-432.
6. Van der Zee FP, Lettinga G, Field JA. The role of (auto) catalysis in the mechanism of an anaerobic azo reduction, *Water Sci Technol* 2000;42:301-308.
7. Mendez-Paz D, Omil F, Lema JM. Anaerobic treatment of azo dye Acid Orange 7 under fed-batch and continuous conditions, *Water Res* 2005;39:771-778.
8. Coates JD, Ellis DJ, Blunt-Harris EL, Gaw C, Roden EE, Lovley DR. Recovery of humic-reducing bacteria from a diversity of environments, *Appl. Environ. Microbiol* 1998;64:1504-1509.
9. Cervantes FJ, de Bok FAM, Tuan DD, Stams AJM, Lettinga G, Field JA. Reduction of humic substances by halorespiring, sulphate-reducing and methanogenic microorganisms, *Environ. Microbiol* 2002;4:51-57.
10. Cervantes FJ, Tuan DD, Ivanova AE, Roest K, Akkermans AD, Lettinga G, Field JA. Selective enrichment of *Geobacter* sulfurreducens from anaerobic granular sludge with quinones as terminal electron acceptors, *Biotechnol Lett* 2003;25:39-45.
11. Hong Y, Xu M, Guo J, Xu Z, Chen X, Sun G. Respiration and growth of *Shewanella* decoloration's S12 with an azo compound as the sole electron acceptor, *Appl Environ Microbiol* 2007;73:64-72.
12. Nie H, Yang F, Zhang X, Yang J, Chen L, Wang J, Xiang Z, *et al.* (11in, Complete genome sequence of *Shigella flexneri* 5b and comparison with *Shigella flexneri* 2a, *BMC Genomics* 2006;7:173.
13. Ghost DK, Mandal A, Chaudhuri J. Purification and partial characterization of two azoreductases from *Shigella dysenteriae* Type 1, *FEMS Microbiol Lett* 1992;77:229-233.
14. Hong Y, Guo J, Xu Z, Xu M, Sun G. Humic substances act as electron acceptor and redox mediator for microbial dissimilatory azoreduction by *Shewanella decolorationis* S12, *J Microbiol Biotechnol* 2007;17:428-437.