



ISSN Print: 2394-7500  
 ISSN Online: 2394-5869  
 Impact Factor: 5.2  
 IJAR 2018; 4(1): 511-512  
[www.allresearchjournal.com](http://www.allresearchjournal.com)  
 Received: 22-11-2017  
 Accepted: 24-12-2017

Virendra Prasad Singh  
 At:-Biraul, Po:-Biraul, Via:-  
 Khajauli, Ps:-Khajauli,  
 Madhubani, Bihar, India

## Synthesis and characterization of substituted acid hydrazides derivative

Virendra Prasad Singh

### Abstract

A novel synthesis and Characterization of benzo derivative of acid hydrazide has been reported. 3-nitro anilines are reacted with chloroacetyl chloride in presence of acetone to give 2-chloro-*N*-(3-nitrophenyl) acetamide which when reacted with 3-amino-3, 4-dihydropyrido [2,3-*b*]pyrazin-2(1*H*)-one in presence of dry pyridine to give the final target. The structures of synthesized compounds are confirmed by their IR and <sup>1</sup>H-NMR spectral data. Melting point of the compound has been determined by open capillary tube.

**Keywords:** acid hydrazide, IR, NMR cyclization, hydrazine.

### Introduction

The important class of heterocyclic is acid hydrazide derivatives. These are reported to possess a wide spectrum of biological properties such as antibacterial [1], analgesic [2], anti-inflammatory [3], antifungal [4], antimalarial [5], antihypertensive [6], CNS depressant [7], anticonvulsant [8], antihistaminic [9], local anesthetic [10], antiparkinsonism [11], anti-viral [12], antitubercular [13], anti-cancer [14] etc. activities.

In this paper a novel synthesis of azocinnoline derivative 1-(3-nitrophenyl)-3-(3-oxo-3, 4-dihydropyrido [2,3-*c*]pyridazin-2(1*H*)-yl)urea has been reported.


### Procedure

13.8 g (or 0.10 mol) of 3-nitroaniline was taken in a round bottom flask to which 50 ml of acetone was added and mixed thoroughly. The 11.2 g (or 0.10 mol) of chloroacetyl chloride was added drop wise to it with continuous shaking. After complete addition, the reaction mixture was refluxed for 3-4 h. The reaction was monitored by TLC. The reaction mixture was cooled and poured into ice-cold water with continuous stirring. Sodium bicarbonate was added to neutralize the hydrogen chloride liberated during the reaction. The product obtained was filtered, thoroughly washed with water, dried and recrystallised with ethanol.

### Synthesis of azocinnoline derivative

#### Procedure

The 0.86 g (0.0031 mol) 2-amino-1,4-dihydropyrido[2,3-*c*]pyridazin-3(2*H*)-one was taken in a round bottom flask and dissolved in 20 ml of dry pyridine then 0.80 g (0.0037 mol) of 2-chloro-*N*-(3-nitrophenyl)acetamide was added and refluxed for 6 h. The reaction was monitored by TLC. After the completion of reaction, the contents were cooled and poured into ice-cold water with continuous stirring and kept aside for 10 min, the crystalline solid obtained was filtered at pump, thoroughly washed with water, dried and recrystallised with ethanol.



Correspondence  
 Virendra Prasad Singh  
 At:-Biraul, Po:-Biraul, Via:-  
 Khajauli, Ps:-Khajauli,  
 Madhubani, Bihar, India

**Table 1:** Mol. Formula, Mol. Weight, Recrystallising Solvent

Mol. formula	Mol. weight	M.P(°C)	Recrystallising solvent	% yield
C <sub>16</sub> H <sub>16</sub> N <sub>4</sub>	264.32	193	Ethanol	52

**References**

1. Martin JN, Rose DA, Hadley WK, Perdreu-Remington F, Lam PK, Gerberding JL. R – J Infect Dis 1999;180:1809-18.
2. Mitsuhashi S. Mol Cell Biochem 1979;26:135.
3. Nikaido H. J Antimicrob Chemother 1988;22:17-22.
4. Tomsaz A, Albino A, Zanati E. Nature 1970;227:138.
5. Chambers HF, Hartman BJ, Tomsaz A. J Clin Invest 1985;76:325-31.
6. Foingold DS. J Med 1969;281:1189.
7. Spera RV, Farber BF. J Amer Med Asso 1992;268:2563-4.
8. Speer BS, Shoemaker NB, Salyers AA. Clin Microbiol, Rev 1992;5:387-99.
9. Burgers Medicinal Chemistry, Part II. 4th ed, New York: Wiley Publishers 1979.
10. Wilson and Giswold's organic, medicinal and pharmaceutical chemistry. 11th ed, Philadelphia: Lippincott Publishers 1998.
11. Kar A. Medicinal Chemistry. 2nd ed. New Age International Publishers, New Delhi 2001.
12. Chambers HF, Hantman BJ, Tomsaz A. J Chin Invest 1985;76:325-31.
13. Wilson and Giswold's Organic, Medicinal and Pharmaceutical Chemistry, 11th ed, Delgado JN, and Remers WA, Eds, Philadelphia, Lippincott Williams and Wilkins 2004;3-271.
14. Korolkovas A. Essentials of medicinal chemistry. 2nd ed, New York: Wiley Publishers 1987.