



ISSN Print: 2394-7500  
ISSN Online: 2394-5869  
Impact Factor: 8.4  
IJAR 2021; 7(10): 280-283  
[www.allresearchjournal.com](http://www.allresearchjournal.com)  
Received: 03-08-2021  
Accepted: 04-09-2021

**Ashok Kumar Namdeo**  
Research Scholar, Department  
of Zoology, Pt. S.N.S. P.G.  
College, Shahdol, Madhya  
Pradesh, India

**Dr. Bharat Sharan Singh**  
Chairman, MP Private  
University Regulatory  
Commission, Bhopal, Madhya  
Pradesh, India

**Corresponding Author:**  
**Ashok Kumar Namdeo**  
Research Scholar, Department  
of Zoology, Pt. S.N.S. P.G.  
College, Shahdol, Madhya  
Pradesh, India

## Physico-Chemical analysis of water of Narmada River at Chandan Ghat Dindori district (M.P.)

**Ashok Kumar Namdeo and Dr. Bharat Sharan Singh**

### Abstract

Water is the basic necessity for a living being and detritions in the quality of water led to many harmful impacts. Water is regarded as 'polluted' when it is changed in its quality or compositions, directly or indirectly as a result of human activities, so that it becomes less suitable for drinking, as well as domestic and other purposes. Pollution of fresh water results largely from the waste disposal. Most of the rivers have become darkened with sewage, chemicals and other undesirable foreign extraneous matter. Moreover, the rivers carry and deposit their pollutants in to the ocean. Hence, the oceans are also polluted by toxic wastes which cause contamination of sea-foods on a large-scale. The present study has been made to analyze the physicochemical parameters of the Chandan ghat, Narmada River. Samples were collected season wise from sampling site for analyzing the various physicochemical parameters such as Temperature, pH, TDS, Conductivity, DO, free CO<sub>2</sub>, Sulphate, Phosphate, Nitrate, BOD, COD. The work highlights the condition of this river water in various seasons with respect to the parameters mentioned above.

**Keywords:** Physico-chemical parameters, Narmada River, Dindori, pollution

### 1. Introduction

Water is a resource that has many uses, including recreation, transportation, and hydroelectric power, domestic, industrial, and commercial uses. Water also supports all forms of life and affects our health, lifestyle, and economic well-being. Although more than three quarters of the earth's surface is made up of water, only 2.8 percent of the Earth's water is available for human consumption (Iskandar, 2010) <sup>[1]</sup>.

Fresh water is a finite resource, essential for agriculture, industry and even human existence, without fresh water of adequate quantity and quality, sustainable development will not be possible (Kumar, 1997) <sup>[2]</sup>. Rivers play a major role in assimilation or carrying off of municipal and industrial wastewater and runoff from agricultural land, the former constitutes the constant polluting source whereas the later is a seasonal phenomenon (Muduli, *et al.* 2010) <sup>[3]</sup>.

A majority of developing countries are in the tropical zone and have fast growing human population. Therefore, there is constant increase in the demand of food, fuel, fiber, medicine and constructions. It results in the exploitation of natural resources. In many countries including India, the rivers are not only being exploited but are also used as dumping places for effluents, sewage and solid wastes. Direct or indirect contact of chemicals or waste water to the sources of drinking water cause the undesirable changes in it which becomes dangerous for all living things. Considerable investigations of physico-chemical properties of the river water are carried out in India (Borse *et al.* 2003, Singh and Gupta 2004, Barai and Kumar 2012, Deshmukh 2012, Chaurasia and Karan 2013, Majumder and Dutta 2014) <sup>[4-9]</sup>.

A water body affects the environment in its vicinity, like charging of ground water tables, conditions of climate etc. Most of the people like washer man, and fisherman, living in the surrounding area depend on this source of water for their survival. Any damages to this water source by any agency will not only make life miserable but that will also disrupt the aquatic ecosystem. It is therefore necessary to study the quality of river water, on the basis of physico-chemical parameters so as to assess its potability.

**2. Material and methods**

A general survey of the river was made for the study of various abiotic parameters. Water has unique property of dissolving and carrying suspension, a huge variety of chemicals, has the undesirable consequence that water can easily become contaminated (APHA 1989) [10]. Water samples were collected monthly in clear glass bottles from surface (max. depth 20 cm) sites of the river. Water samples were collected in three replicates from surface, column and bottom and mean values of all three observations were taken into consideration. For BOD estimation, water samples were collected separately in dark bottles. The acquisition of meaningful data demands correct sampling and storage procedures. The preservation of samples were done by refrigeration at 4°C, which is most general accepted method. Water and air temperature were recorded with a digital centigrade thermometer on the date of sampling.

Physico-chemical parameters like water temperature, pH, DO, free Carbon-di-oxide, total alkalinity and conductivity were measured in the field. Other parameters were mostly

tested within 24 hrs of collection. A total of 12 limnological parameters of water viz., temperature, pH, DO, BOD, COD, Free CO<sub>2</sub>, total alkalinity, conductivity, TDS, Phosphates, nitrate were determined. All the parameters were analysed the standard methods (Golterman 1969, Michael 1984, Trivedy and Goel 1986 and APHA 1989) [11, 12, 13, 10] and spectrophotometer SQ 118.

**3. Result and Discussion**

The present study was conducted at selected sampling station of Chandan ghat, Narmada River at Dindori district for a period of one year (from Jan 2020 to Dec. 2020). Covering three main seasons i.e. Rainy (July/August/September), Winter (Dec/Jan/Feb) and Summer (Apr/May/June) in a year.

Physico-chemical and bacteriological parameters were carried out in the samples collected from the study area to study the drinking water quality and pollution level and details of the same was given in table-1 to show the seasonal fluctuations of selected parameters (table-1).


**Table 1:** Physico-chemical properties of Chandan ghat, Narmada River for the duration Jan. 2020-Dec.2020.

S. No.	Parameters	Rainy Seasons	Winter seasons	Summer seasons	Mean	SD
1.	Temp. (°C)	28.9	21.2	30.4	26.83	4.94
2.	Transperancy (cm.)	91	112	86	96.33	13.80
3.	T.D.S. (mg/l)	379	283	321	327.67	48.35
4.	Conductivity	374	275	411	353.33	70.32
5.	pH	7.5	8.6	7.9	8.00	0.56
6.	Alkalinity (mg/l)	136	174	196	168.67	30.35
7.	D.O. (mg/l)	7.6	9.2	5.15	7.32	2.04
8.	B.O.D. (mg/l)	13.9	9.16	29.8	17.62	10.81
9.	C.O.D. (mg/l)	119.4	139.8	82.2	113.80	29.21
10.	Nitrates (mg/l)	0.029	0.022	0.023	0.02	0.00
11.	Phosphates (mg/l)	0.049	0.067	0.071	0.06	0.01
12.	Sulphate	17.3	12.9	18.91	16.37	3.11

To assess the quality of river, Indian drinking water quality standard IS 10500 (1990) has been adopted. The data harvested during the present study is given in table 1. The present data showed the seasonal variations of all the parameters during the study period.

The water temperature is one of the important parameter in river. In the present study of Narmada River, difference in

the fluctuation of water temperature was maximum 30.4°C to 21.2°C with a standard deviation of 4.94. The season wise studies showed the increased temperature of the river during summer is due to the common effect of intensity of solar radiations, ambient temperature. Decreased values of temperature during winter days are due to low ambient temperature (Gyananath *et al.* 2000) [14].



**Fig. 1 :** Graphics Average water analysis of Chandan ghat, Narmada river.

**Fig 1:** Graphics Average water analysis of Chandan ghat, Narmada River

Transparency is light penetration capacity of the water. The color of water is due to concentration of suspended organic and inorganic particles. Transparency varied from maximum 112 cm to 86 cm. with a standard deviation of 13.80. The less transparency observed during rainy season and summer season. While during winter comparatively the water showed more transparency. Low transparency in summer and rainy season in the different water bodies in Jammu (Shashi and Raina 1990, Kaushik and Saksena 1999 and Sharma 2015) [15-17].

Total dissolved solids include salt and variety of organic substances, which readily dissolve in water and often impart a degree of hardness. The value of total dissolved solids ranged between max 379 mg/ltr to min 283 mg/ltr with a standard deviation of 48.35. The maximum seasonal value is observed in rainy season and minimum value in winter season. Moreover the low value of total dissolved solids in summer and high in rainy season was also observed (Trivedy *et al.* 1984) [18].

The ionic status of water determines the conductivity. During summer and rainy days the enhanced values of conductivity may be due to presence of carbonates and bicarbonates in the water in more quantity due to pollutants. The exchange of inorganic ions by increased micro flora during the winter showed the moderate values of conductivity during the winter period.

Hydrogen ion concentration is considered as a important ecological factor, which is a result of interaction of various substances in water and in numerous biological phenomenon. Nearly neutral pH of water is regulated by carbon dioxide and bicarbonates (Hutchinson, 1957) [19]. The river water showed well alkaline water through the study period. pH of river ranges between 7.5 to 8.6.

Total alkalinity in river water ranges from 136mg/l to 196mg/l with a standard deviation of 30.35. During the summer season the higher values of hardness were observed and lower values during winter season (Ugale and Hiware 1999, Pratibha *et al.* 2005 and Sharma 2015) [20-21, 17].

Dissolved oxygen is also one of the important factors of water quality, which influences the biota present inside the river water. The seasonal fluctuation of dissolved oxygen in water bodies (Adebisi 1981) [22]. Similar pattern of DO observed in the present study, it decreases during the hot

days of summer. DO show the inverse relationship with the temperature in river water.

Biological oxygen Demand is a direct measure of O<sub>2</sub> requirement and indirect measure of biodegradable organic matter. The maximum B.O.D. was 29.8mg/l in summer and minimum 9.16mg/l in winter season. (Seenaya and Zafar 1979 & Sharma 2015) [23, 15].

Chemical Oxygen Demand indicates the extent of chemical pollution mainly from industrial effluents. The C.O.D. values observed maximum in winter and minimum in rainy season (Singh and Roy, 1995 & Sharma, 2015) [24, 15]. In the present study, nitrate values ranged between 0.029 to 0.022 mg/l maximum of nitrate values were reported during rainy season and summer and minimum during winter. The high value of nitrate in rainy season, which is linked to heavy run-off of the organic matter from the catchments (Gohram, 1961) [25].

Phosphate concentration in river water ranged between 0.071mg/l to 0.049mg/l. more concentration of phosphates recorded during the summer and winter season. It may be due to deposition of ashes and bones under religious activities and decomposition of organic matter in the water sediments.

Sulphate is produced by biological oxidation of sulphur content of organic matter. The sulphate value ranges between 18.91mg/l to 12.9mg/l with a standard deviation of 3.11. Minimum concentration of sulphate was in winter and maximum in summer (Angadi *et al.* 2005 & Sharma 2015) [26, 15].

#### Test of significance of the observed correlation coefficient

The correlation coefficient study is very useful to determine a predictable relationship which can be exploited in practice. It is used for the measurement of the strength and statistical significance of the relation between two or more water quality parameters. Hence, it is a helpful tool for the promotion of research activities (Shrivastava and Joshi, 2008 and Borkar and Tembhre, 2018) [27-28]. The correlation coefficients (r) among the various physicochemical characteristics of Effluents from recycled Orient paper mill, Amlai have been calculated and the numerical values are tabulated as shown in Table-2.

**Table 2:** Correlation Matrix of Physico-chemical properties of Chandan ghat, Narmada River for the duration Jan. 2020-Dec. 2020

S.No.	Temp.	Transparency	T.D.S.	Conductivity	pH	Alkalinity	D.O.	B.O.D.	C.O.D.	Nitrates	Phosphate	Sulphate
Temp.	1											
Trans-perancy	-1.00	1										
T.D.S.	0.70	-0.68	1									
Cond-uctivity	0.99	-1.00	0.61	1								
pH	-0.87	0.85	-0.96	-0.81	1							
Alkal- inity	0.00	-0.03	-0.71	0.11	0.50	1						
D.O.	-0.88	0.90	-0.28	-0.93	0.53	-0.47	1					
B.O.D.	0.78	-0.80	0.10	0.85	-0.37	0.62	-0.98	1				
C.O.D.	-0.86	0.87	-0.23	-0.91	0.49	-0.51	1.00	-0.99	1			
Nitrates	0.48	-0.46	0.96	0.38	-0.85	-0.88	-0.01	-0.17	0.03	1		
Phos-phates	-0.20	0.17	-0.84	-0.09	0.66	0.98	-0.29	0.46	-0.33	-0.95	1	
Sulphate	0.99	-1.00	0.62	1.00	-0.81	0.11	-0.93	0.84	-0.91	0.38	-0.09	1

In the Correlation Matrix of Physico-chemical properties of Chandan ghat, Narmada river, out of 78 correlation coefficients, 36 negative and 42 positive correlation coefficients. In Table-2, the highly positive correlation is observed between Sulphate and conductivity (1.00), Conductivity and Temperature (0.99), Sulphate and

Temperature (0.99), Phosphate and alkalinity (0.98), Nitrates and TDS (0.96) and DO and Transparency (0.90), where highly negative correlation is observed between Transparency and Temperature (-1.00), Conductivity and Transparency (-1.00), COD and BOD (-0.99), BOD and DO

(-0.98), Phosphate and Nitrate (-0.95), DO and Conductivity (-0.93) and COD and Conductivity (-0.91).

#### 4. Conclusion

The objective of this research work was to find the quality of Chandan ghat, Narmada river water w.r.t drinking purpose. From the experimental results of water quality analysis of Narmada River, it can be concluded that the water quality is good and most of the parameters are within the limits set by organizations like WHO and Bureau of Indian Standard (BIS). From the above study, it may conclude that except little variation, all the physico-chemical parameters were in permissible limit at the study site of the Chandan Ghat, Narmada River. It is suggested that proper measures are necessary to avoid contamination as water is used for drinking purpose. At present the river is suitable for irrigation and fishery purpose.

#### 5. Acknowledgement

The authors are greatly indebted to Principal and Head of Zoology Dept. of Pt. S.N.S. Govt. Science P.G. College, Shahdol (M.P.) who permitted to carry out this work.

#### 6. References

1. Iskandar MB. The effectiveness of bio-filter as a treatment for domestic wastewater, University Malaysia Pahang (thesis) 2010.
2. Kumar NA. View on Freshwater environment, Ecol. Env and Cons 1997, 3(3).
3. Muduli Bipra Prasanna, Panda Chitta Ranjan. Physico chemical properties of water collected from Dhamra estuary, International journal of environmental sciences 2010, 1(3).
4. Borse SK, Lohar PS, Bhawe PV. Hydrobiological study of algae of Aner river, Jalgaon (Maharashtra), J Aqu. Bio 2003;18(1);15-18.
5. Singh M, Gupta KC. Physico-chemical studies of water of river Yamuna at Mathura, Ecol. Envi. And Cons 2004;10(2):193-196.
6. Barai SR, Kumar Satish. Evaluation of the physicochemical characteristics of river Varuna at Varanasi, India, J Environ. Biol 2012;34:259-265.
7. Deshmukh BS. Hydrobiological study of river Pravara in Ahmednagar district (Maharashtra), Bionano Frontier, Eco Revolution 2012 Colombo 2012, 89-92.
8. Chaurasia S, Karan Raj. Water quality and pollution load of river Mandakini at Chitrakoot, India, Int. Res. J Environ. Sci 2013;2(6):13-19.
9. Majumder S, Dutta TK. Studies on seasonal variations in physico-chemical parameters in Bankura segment of the Dwarakeshwar River (W.B) India, IJAR 2014;2(3):877-881.
10. APHA. Standard methods for the examination of water and wastewater. 17th edition, American Public Health Association, Washington D.C 1989.
11. Golterman HL (Ed). Methods of chemical analysis of freshwaters. IBP Handbook No. 8, Blackwell, Oxford 1969.
12. Michael P. Ecological methods for field and laboratory investigations. Tata-McGraw Hill Pub. Com. Ltd., New Delhi 1984, 404.
13. Trivedy PK, Goel RK. Chemical and Biological methods water pollution studies, Environmental publication Karad India 1986.
14. Gyananath G, Shevnikar SV, Samiuddin S. Water quality analysis of river Godavari during Holimela at Nanded, Poll. Res 2000;19(4):673.
15. Shashi Kant, Raina AK. Limnological studies of two ponds in Jammu. J Env. Biol 1990;11(2):137-144.
16. Kaushik S, Saksena DN. Physico-chemical limnology of certain water bodies of central India. In: fresh water ecosystems of India (Ed. K. Vijay Kumar) Daya Publ., House, Delhi 1999, 1-58.
17. Sharma Pankaja. Seasonal Variations in Physico-Chemical Properties of Narmada River in Dindori Madhya Pradesh, India, International Journal for Research in Applied Science & Engineering Technology (IJRASET), Volume 3 Issue XII, December 2015, 285-288.
18. Trivedy RK, Goel PK, Shrotri AC, Khatavkar SD. Prospective in Aquaculture, Biol 1984, 15-18.
19. Hutchinson GE. A treatise on limnology vol. II John Wiley and Sons, New York 1957, 1015.
20. Ugale BJ, Hiware CJ. Limnological study of an ancient reservoir Jagtunga Samudra located at Kandhar, Dist. Nanded, Maharashtra, India. Eco. Envi and Cons 1999;11(3-4):473-475.
21. Pratibha V, Raithak, Bhuktar AS. Physicochemical analysis of drinking water from different areas of Aurangabad, Poll. Res 2005;24(3):727-728.
22. Adebisi. The physicochemical hydrobiology of a tropical river upper Ogun river Nigeria. Hydrobiol 1981;79(2):157-165.
23. Seenaya G, Zafar AR. An ecological study of Mir Alam Lake, Hyderabad, India, Indian J Biol 1979;11(2):330-335.
24. Singh JP, Roy SP. Limnobiological investigation of Karwarlake, Begusarai, Bihar Env. Eco 1995;13:330-335.
25. Gohram. The chemical composition of some waters from Dune slacks at Sandscale, North Lancshire, J Ecology 1961;49(1):79-82.
26. Angadi SB, Shiddamallayya N, Patil PC. Limnological studies of Papanash pond, Bidar, Karnataka, J Env. Biol 2005;26(2):213-216.
27. Borkar Pranjali, Tembhe Manju. Comparative analysis of physico-chemical properties of water of five lakes of Bhopal, India, World Journal of Pharmaceutical Research 2018;7(7):2228-2242.
28. Shrivastava K, Joshi S. Physico-chemical investigation and correlation analysis of water quality of Upper Lake of Bhopal, M.P., (India). Curr World Environ 2008;3(2):327-330.