



ISSN Print: 2394-7500
ISSN Online: 2394-5869
Impact Factor: 8.4
IJAR 2022; 8(10): 111-114
www.allresearchjournal.com
Received: 12-07-2022
Accepted: 09-09-2022

Rajesh Kumar Mishra
Research Scholar Botany,
Govt. Science P.G. College,
Rewa, Madhya Pradesh, India

Santosh Kumar Agnihotri
Professor of Botany, Govt.
Science P.G. College, Rewa,
Madhya Pradesh, India

Ecological studies of tree vegetation of Chanda forest range of Dindori district (M.P.) India

Rajesh Kumar Mishra and Santosh Kumar Agnihotri

Abstract

Ecological studies are the measure of biodiversity status of any vegetation. Knowledge of ecological composition, diversity of tree species and dominant communities of the ecosystem is the basis of planning and implementation of biodiversity conservation management. To fulfil this requirement the study was carried out at Chanda forest of Dindori District, Madhya Pradesh, India. The study provided the composition of tree communities and the status of biodiversity of these trees. The article mainly focused on the diversity and ecology of the tree vegetation of the forest range. A total of 21 established tree species belonging to 17 families were recorded. The present studies were made to evaluate the Frequency, Density, Basal Area and Importance Value Index along with some community indices.

Keywords: Biodiversity, quadrat, phytosociology, dominance, co-ordinance, basal area, IVI

Introduction

Forests are the greatest resources among the ecosystems that support life throughout the globe. The forest ecosystems are the most rich terrestrial ecosystems which are characterized by the high species diversity and species richness. These ecosystems are distinguished from all other terrestrial ecosystems by a very high level of biodiversity. Now a days biodiversity is declining seriously on a global scale, which also reveals the importance of conservation planning. Except protected areas and reserve forests, scientific studies on biodiversity of special habitats in India are meager like the other areas of the world. Furthermore, extensive surveys of biodiversity have been conducted only for a few taxonomic groups and ecosystem types. The situation is the same in Chanda range of Dindori district. Only preliminary information on the biodiversity patterns of plant groups are available in this forest range. Such a lack of study significantly hinders the assessment of the value of existing species, their current status and threats which might facilitate their long term conservation (Lohbeck *et al.* 2014) ^[1]. As a result, a considerable number of species may currently be under threat of local extinction. From the ecological point of view, eliminating or decreasing forest ecosystem has severe negative effect on other ecosystems (Palit *et al.* 2012) ^[2]. The present investigation has been conducted to assess the status of biodiversity of tree species in Chanda Forest Range.

For the evaluation of the status several phytosociological and quantitative parameters were determined. Such quantitative description of plant patterns provides a clear view to look at the interaction between the plants as well as between the plant and environment. It can also provide a clearcut evidence about the status of biodiversity (Chase and Leibold, 2003) ^[3].

Materials and Methods

Description of Study Site: Chanda forest range is expanded in eastern part of the Dindori district. It is 56 km. away from district headquarter and situated in Maikal hills at the side of Jabalpur-Amarkantak National highway. The height of the forest is 885 ft. minimum from msl and 1100 ft. maximum from msl.

In the east of Chanda forest, there is Karanjia range, in west & south. Samnapur range and in north Gadasarai range are situated. The Chanda Forest range has expanded between 22° to 22.50° North latitude and 81.15 to 81.20 longitude. The present study area has been divided into 3 sites *viz.* Chanda forest, Tantar forest, and Silpidi forest.

Corresponding Author:
Rajesh Kumar Mishra
Research Scholar Botany,
Govt. Science P.G. College,
Rewa, Madhya Pradesh, India

Methodology: For phytosociological studies in Chanda forest range, the quadrat methods were used. In each forest beat five quadrates laid down for trees. The sizes of quadrates for trees were 10 m.sq. Basal area was calculated from the perimeter which was measured at a breast height (Phillips, 1959) [6].

Data Analysis Techniques: To analyses the level of diversity in tree vegetation several phytosociological parameters like frequency, Relative frequency, density and Relative density etc., were calculated (Phillips, 1959; Chaubey *et al.* 1988 and Misra, 1968) [6-8]. Then IVI of trees were made to determine the dominant species of the forest. Dominance is a significant indicator of species composition in a forest ecosystem (Burak *et al.* 2011; Sahu *et al.* 2008) [9,10]. The dominance of any species refers to its relative value or importance in its habitat (Chase and Leibold, 2003) [3]. Or in other language it is the measure of the degree of influence of the species on the ecosystem. To assess the over all impact of a species Importance Value Index was determined by adding Relative frequency, Relative density and Relative Basal Area (Misra, 1968 and Priya *et al.* 2005) [8, 11].

Frequency (%): Frequency refers to the degree of dispersion of individual species in an area and usually expressed in terms of percentage. It is calculated by the equation:

$$\text{Frequency (\%)} = \frac{\text{No. of plot in which the species is present}}{\text{Total No. of plots sampled}} \times 100$$

Density: Density refers to the expression of the numerical strength of a species. It is calculated by the equation:

$$\text{Density} = \frac{\text{No. individuals of the species}}{\text{Total No. of plots sampled}}$$

Relative Frequency (%): Relative Frequency is the degree of dispersion of individual species in an area in relation to the number of all the species occurred.

$$\text{Relative Frequency (\%)} = \frac{\text{Frequency of the species}}{\text{Frequency of all the species}} \times 100$$

Relative Density (%): Relative Density is the measure of numerical strength of a species in respect to the total number of individual of all the species. It can be determined by the equation.

$$\text{Relative Density} = \frac{\text{Density of the species}}{\text{Density of all the species}}$$

Relative Dominance (%): Dominance is the parameter which is determined by the value of basal area. For the comparative analysis Relative dominance is determined. It is the coverage value of a species with respect to the sum of coverage of the rest of the species in the area.

$$\text{Basal area} = \frac{(\text{Circumference at breast height})^2}{12.56}$$

Relative dominance or Relative Basal Area

$$= \frac{\text{Basal Area of the species}}{\text{Basal area of all the species}}$$

Importance Value Index: Importance Value Index is used to determine the over all impact of each species in the community structure. It is calculated by the addition of the percentage values of the relative frequency, relative density and relative dominance (Relative Basal Area).

$$\text{IVI} = \text{Relative Frequency} + \text{Relative Density} + \text{Relative}$$

Data Processing and Phytosociological Analysis: All the phytosociological data collected from different sources were tabulated and analysed individually. The data collected were used to compute some community indices like.

Species diversity (H'): Species diversity was determined by the Shannon-Weiner Index (Shannon and Wiener, 1963) [12]. It was calculated by the equation,

$$(H') = - \sum [(ni/N) \cdot \ln (ni/N)]$$

Where ni= IVI of individual species and N= total IVI of all the species (Shannon and Wiener, 1963) [12].

Species dominance (Cd): Species dominance was calculated by the Simpson Index (Simpson, 1949) [13]: Cd = $\sum (ni/N)^2$,

Where ni= IVI of individual species and N= total IVI of all the species.

Equitability of evenness (e): Equitability of evenness is the measure of the degree of relative dominance of each species in the habitat. It was determined according to Pielou (1966) [14] as:

$$\text{Evenness (e)} = H'/\log S$$

Where: H'= Shannon index, S = number of species.

Species richness (D): Species richness was calculated by Margalef (1968) [15] Index as:

$$D = (S-1)/\ln N.$$

Where: S = number of species. N= total number of individuals

Menhinick's index (D_{mm}): Menhinick's index (Whittaker and Levin, 1977) [16] is expressed as

$$D_{mm} = S/N,$$

Where: N= Number of individuals in the sample, S= Number of species.

Equitability Index: The Shannon's equitability Index (Simpson, 1949) [13] is expressed as

$$(EH) = H'/H_{\max} = H'/\ln S$$

Berger-Parker Dominance Index: The Berger-Parker

Dominance Index is the measure of numerical importance of the most abundant species. It is determined by the equation $d = N_{\max}/N$.

Where: N_{max} = Number of individuals of the most abundant species, N = Total number of individuals in the site.

Results and Discussion

Observation of this study indicates that *Adina cordifolia* Hook.f. had highest density (3.80) and *Shorea robusta* Gaertn. f. had maximum IVI (48.65). Density of *Shorea robusta* Gaertn. f. was recorded as 2.80. *Adina cordifolia* Hook.f. had IVI value as 30.49. It is also noted that some other tree species had a good IVI value. These include *Wrightia tomentosa* Roem and Schult. (23.39), *Lagerstromia perviflora* Roxb. (17.92) and *Butea monosperma* Taub. (17.04). However maximum relative basal growth is recorded for *Shorea robusta* Gaertn. f. and it is 4.35. Relative basal growth is not so good in other tree species of this forest including *Adina cordifolia* Hook.f. (4.35), *Wrightia tomentosa* Roem. and Schult. (3.63) and *Butea monosperma* Taub. (0.92). Relative basal area is also found good for *Sterculia villosa* Roxb. (13.1). In this survey it is also recorded that few species had very low IVI and

Relative basal area *Albizia procera* has minimum IVI value of 1.80 and relative basal area of 0.34.

To assess the overall status of biodiversity in Chanda Forest Range, eight diversity indices were used. Shannon and Weiner (1963) [12] index represents entropy. It is a diversity index that consider the number of individual species as well as the number of taxa. It ranges from zero to higher value. The communities with only single taxa has the value of zero. Increase of the value of diversity index reveal higher number of taxa in the community. Simpson's dominance Index was also found much less than 1, which showed that the sites were not dominated by single species (Huston, 1994) [18]. On the contrary a few species dominate the forest. The primary conclusion is that there is low grazing pressure and moderate human impact on normal distribution of tree species which may cause reduction in tree community in next few decades in the forest ecosystem. Both the Menhinick's Index and Margalef's Index measure richness of species in the ecosystem.

Table 1: Value for the different phytosociological parameters measured for different tree species of Chanda forest range

Name of the Plant	Family	A	D	Fr (%)	BA	RD	RF	RBA	IVI
<i>Adina cordifolia</i> Hook.f.	Rubiaceae	4.22	3.80	90	810.29	16.96	9.18	4.35	30.49
<i>Aegle marmelos</i> Correa	Rutaceae	1.75	0.70	40	718.55	3.12	4.08	3.86	11.06
<i>Albizia procera</i> (Roxb.) Benth. gum	Fabaceae	1.00	0.10	10	64.66	0.44	1.02	0.34	1.8
<i>Amoora walichii</i> King.	Meliaceae	1.80	1.00	50	1391.95	4.46	5.1	7.48	17.04
<i>Anthocephalus cadamba</i> Miq.	Rubiaceae	1.50	0.30	30	247.72	1.33	3.06	1.33	5.72
<i>Bischofia javanica</i> Blume	Phyllanthaceae	2.66	1.60	60	473.49	7.14	6.12	2.54	15.8
<i>Butea monosperma</i> (Lamk.) Taub	Fabaceae	1.50	0.60	40	171.14	2.67	4.08	0.92	7.67
<i>Careya arborea</i> Roxb.	Lecythidaceae	2.16	1.30	60	210.32	5.8	6.12	1.13	13.05
<i>Dillenia pentagyna</i> Roxb.	Dilleniaceae	1.71	1.20	70	684.91	5.35	7.14	3.68	16.17
<i>Lagerstromia perviflora</i> Roxb.	Lythraceae	2.00	1.20	60	1199.73	5.35	6.12	6.45	17.92
<i>Lagerstromia speciosa</i> (L.) Pers.	Lythraceae	2.00	0.20	10	33.16	0.89	1.02	0.17	2.08
<i>Madhuca indica</i> Gmel.	Sapotaceae	1.50	0.03	30	678.76	1.33	3.06	3.65	8.04
<i>Shorea robusta</i> Gaertn. f.	Dipterocarpaceae	2.80	2.80	100	4825.83	12.5	10.2	25.95	48.65
<i>Sterculia villosa</i> Roxb.	Sterculiaceae	2.00	0.20	10	2436.30	0.89	1.02	13.1	15.01
<i>Syzygium cumini</i> (Linn.) Skeels	Myrtaceae	1.33	0.40	10	392.53	1.78	1.02	2.11	4.91
<i>Tectona grandis</i> Linn.	Verbenaceae	1.25	0.50	40	170.15	2.23	4.08	0.91	7.22
<i>Terminalia alata</i> Heyne ex Roth.	Combretaceae	2.25	0.90	40	595.39	4.01	4.08	3.2	11.29
<i>Terminalia bellerica</i> (Gaertn.) Roxb.	Combretaceae	2.20	1.10	50	554.26	4.91	5.1	2.98	12.99
<i>Terminalia tomentosa</i> Roth.	Combretaceae	1.75	1.40	80	435.62	6.25	8.16	2.34	16.75
<i>Toona ciliata</i> Roem.	Meliaceae	1.00	0.20	20	1819.78	0.89	2.04	9.78	12.71
<i>Wrightia tomentosa</i> Roem.& Schult.	Apocynaceae	3.25	2.60	80	676.20	11.6	8.16	3.63	23.39

Table 2: Value for the different community index parameters, measured for different tree species of Chanda forest range

Sl. No.	Name of the Plant	Shannon Index (H)	Species dominance	Evenness	A/F index
1	<i>Adina cordifolia</i> Hook.f.	0.2331	0.0104	0.1763	0.046
2	<i>Aegle marmelos</i> Correa	0.1196	0.0012	0.0904	0.043
3	<i>Albizia procera</i> (Roxb.) Benth. gum	0.0306	0.00003	0.0231	0.100
4	<i>Amoora walichii</i> King.	0.1630	0.0032	0.1232	0.036
5	<i>Anthocephalus cadamba</i> Miq.	0.0756	0.0003	0.0571	0.050
6	<i>Bischofia javanica</i> Blume	0.1550	0.0027	0.1172	0.044
7	<i>Butea monosperma</i> (Lamk.) Taub	0.0943	0.0006	0.0713	0.037
8	<i>Careya arborea</i> Roxb.	0.1363	0.0019	0.1031	0.036
9	<i>Dillenia pentagyna</i> Roxb.	0.1583	0.0029	0.1197	0.024
10	<i>Lagerstromia perviflora</i> Roxb.	0.1682	0.0035	0.1272	0.033
11	<i>Lagerstromia speciosa</i> (L.) Pers.	0.0343	0.00004	0.0259	0.200
12	<i>Madhuca indica</i> Gmel.	0.0969	0.0007	0.0730	0.050
13	<i>Shorea robusta</i> Gaertn. f.	0.2948	0.2948	0.2229	0.028
14	<i>Sterculia villosa</i> Roxb.	0.1497	0.0025	0.1132	0.200
15	<i>Syzygium cumini</i> (Linn.) Skeels	0.0670	0.0002	0.0506	0.133
16	<i>Tectona grandis</i> Linn.	0.0895	0.0005	0.0624	0.031
17	<i>Terminalia alata</i> Heyne ex Roth.	0.1233	0.0014	0.0932	0.056
18	<i>Terminalia bellerica</i> (Gaertn.) Roxb.	0.1359	0.0018	0.1028	0.044
19	<i>Terminalia tomentosa</i> Roth.	0.1595	0.0030	0.1206	0.021
20	<i>Toona ciliata</i> Roem.	0.1331	0.0017	0.1006	0.050
21	<i>Wrightia tomentosa</i> Roem.& Schult.	0.1989	0.0060	0.1989	0.040

Table 3: Value for different community indices for Chanda Forest Range

Community indices	Value
Species diversity (H')	2.827
Species dominance (Cd)	0.3394
Equitability of evenness (e)	2.129
Species richness (d)	3.695
Menhinick's index (D _{mm})	0.0938
Equitability Index	0.924
Berger-Parker Dominance Index	0.176

Conclusion

The paper reflects the phytosociological characters of tree vegetation of Chanda forest range in Dindori forest division, Madhya Pradesh, India. This study implies the variety of tree species, their distribution and the status of dominance. The vegetation of the Chanda forest range is composed of mosses, ferns, native grasses, sedges, climbers, shrubs and trees. It is also rich in faunal diversity. Here Diversity index of tree species was found as 2.827, whereas dominance index (Cd) was observed as 0.3394. Both the indices reflect that the forest patch is rich in tree vegetation and tree diversity. The present investigation also revealed some interesting phytosociological findings about the tree vegetation of the forest. The findings have illustrated that most abundant plant species *i.e.*, *Adina cordifolia* Hook.f. Lack proper growth (growth of basal area), whereas species having highest basal growth *i.e.*, *Adina cordifolia* Hook.f. are comparatively less abundant. In addition another two species, *Wrightia tomentosa* Roem. and Schult and *Bischofia javanica* Blume are also abundant. Another noticeable fact is that IVI of *Wrightia tomentosa* Roem. and Schult is more than the IVI of *Shorea robusta* Gaertn. f. Thus more than one species are dominant in this forest. This finding supports theories of co-dominant succession.

Therefore, the study recommends further research to be carried out to study succession pattern including tree species loss more specifically (*i.e.* effects of specific species in specific ecosystems such as in forest, grassland and bush-land ecosystems) regeneration ability of the previously disturbed species in this forest range.

Acknowledgement

The authors are thankful to authority of forest department of Dindori district and research centre of Govt. Science P.G. College, Rewa (M.P.) India for giving permission and providing necessary facilities.

References

- Lohbeck M, Poorter M, Martinez-Ramos J, Rodriguez Velazquez M, Van Breugel, Bongers F. Changing Drivers of Species Dominance during Tropical Forest Succession. *Functional Ecology*. 2014;28(4):1052-1058.
- Palit D, Pal S, Chanda S. Diversity and richness of plants in Darjeeling Himalaya with an eye on Gaddikhana forest beat, Senchal east zone forest range, Darjeeling. *Indian Journal of Forestry*. 2012;35(1):39-44.
- Chase JM, Leibold MA. *Ecological Niches: Linking Classical and Contemporary Approaches*. University of Chicago Press, Chicago; c2003.
- Rannie WF. Summer air temperature and number of vascular species in arctic Canada. *Arctic*. 1986;39:133-137.
- Warman CK. *Trees of India (Medicinal, Commercial, Religious & Ornamental) (A Colour Atlas)*. S.K. Jain for CBS Publisher & Distributors, Daryaganj, New Delhi; c1999.
- Phillips EA. *Methods of vegetation study*. Henri Holt Co. Inc; c1959.
- Chaubey OP, Prasad R, Mishra GP. Studies of Teak Plantation and mixed natural forest in Madhya Pradesh. I. Phytosociology, distribution, species diversity and quantitative parameters of tree species *J. Trop. For.* 1988;4:22-35.
- Misra R. *Ecology Work Book*. Oxford & IBH Publishing Co; c1968.
- Burak KP, Roy W, Matthias MB, Craig M, Pauline FG. Response of Plant Species and Life Form Diversity to Variable Fire Histories and Biomass in the Jarrah Forest of South-West Australia. *Austral Ecology*. 2011;37(3):330-338.
- Sahu PK, Sagar R, Singh JS. Tropical Forest Structure and Diversity in relation to altitude and disturbances in a Biosphere reserve in Central India. *Applied Vegetation Science*. 2008;11(4):461-470.
- Priya D, Jean PP, Egbert GL. Changes in Rain Forest Tree Diversity, Dominance and Rarity across a Seasonality Gradient in the Western Ghats, India. *Indian Journal of Biogeography*. 2005;32(3):493-501.
- Shannon CE, Wiener W. *A mathematical theory of Communication*. University Illinois Press, Urban; c1963.
- Simpson EH. Measurement of Diversity. *Nature*, 1949;163(4148):688.
- Pielou EC. Species Diversity and pattern diversity of in the study of Ecological; Succession. *Journal of theoretical Biology*. 1966;10(2):370-383.
- Margalef R. *Perspective in Ecological Theory*. Uni. of Chicago Press; c1968.
- Whittaker RH, Levin SA. The role of mosaic phenomena in mosaic communities. *Theoretical Population Biology*. 1977;12(2):117-139.
- Berger WH, Parker FL. Diversity of plank tonic Foraminifer in deep sea Sediments *Science*. 1970;168(3937):1345-1347.
- Huston MA. *Biological Diversity: the coexistence of species*. Cambridge University Press, Cambridge; c1994.