



ISSN Print: 2394-7500
ISSN Online: 2394-5869
Impact Factor: 8.4
IJAR 2023; 9(5): 106-107
www.allresearchjournal.com
Received: 10-01-2023
Accepted: 11-02-2023

Dr. Shaheen Hameed
Postgraduate,
Department of Orthopaedics,
Sree Mookambika Institute of
Medical Sciences, Kulasekharam,
Kanyakumari, Tamil Nadu,
India

Dr. Uvarajan T
Postgraduate,
Department of Orthopaedics,
Sree Mookambika Institute of
Medical Sciences, Kulasekharam,
Kanyakumari, Tamil Nadu,
India

Dr. Sukaash AJ
Postgraduate,
Department of Orthopaedics,
Sree Mookambika Institute of
Medical Sciences, Kulasekharam,
Kanyakumari, Tamil Nadu,
India

Dr. Binesh SL
Postgraduate,
Department of Orthopaedics,
Sree Mookambika Institute of
Medical Sciences, Kulasekharam,
Kanyakumari, Tamil Nadu,
India

Corresponding Author:
Dr. Shaheen Hameed
Postgraduate,
Department of Orthopaedics,
Sree Mookambika Institute of
Medical Sciences, Kulasekharam,
Kanyakumari, Tamil Nadu,
India

The effectiveness of corticosteroid injection in the treatment of tennis elbow

Dr. Shaheen Hameed, Dr. Uvarajan T, Dr. Sukaash AJ and Dr. Binesh SL

Abstract

Introduction: Lateral Epicondylitis, also known as "Tennis Elbow" is the most common overuse syndrome in the elbow. It is a tendinopathy injury involving the extensor muscles of the forearm. These muscles originate on the lateral epicondylar region of the distal humerus. Tennis elbow affects 1 - 3% of general population. Lateral epicondylitis can occur during activities that require repeated supination and pronation of the forearm with the elbow in near full extension. Most of patients with tennis elbow can be managed conservatively; treatment is successful in 95% of patients.

Materials and Methods: The present study is the effectiveness of local corticosteroid injection in the management of "tennis elbow" was conducted in postgraduate Department of Orthopaedics, Sree Mookambika institute of medical sciences from September 2020 to August 2021. 60 patients with tennis elbow of either sex were included in the study. After taking written informed consent all the patients received local steroid injection (Methylprednisolone 40 mg mixed with 1 ml injection of lignocaine 2%)

Results: Out of 60 patients in our study 41 (68.3%) patients obtained complete relief of pain at six weeks' time. But at final follow-up of 12 weeks only 14 (24.35%) patients were pain free. At 12 weeks follow-up 40% (24) of patients had recurrence of symptoms.

Conclusion: In patients with tennis elbow use of methylprednisolone gave good response for short term use and showed to be an excellent treatment. This procedure is not expensive, mini-invasive, with low surgical risk and short hospitalization.

Keywords: Tennis elbow, corticosteroid injection, inflammation, Methylprednisolone

1. Introduction

Tennis elbow, or lateral epicondylitis, is a painful condition of the elbow caused by overuse. Playing tennis or other racquet sports can cause this condition. Tennis elbow is inflammation or, in some cases, microtearing of the tendons that join the forearm muscles on the outside of the elbow. The forearm muscles and tendons become damaged from overuse - repeating the same motions again and again. This leads to pain and tenderness on the outside of the elbow. There are many treatment options for tennis elbow. In most cases, treatment involves a team approach. Primary doctors, physical therapists and, in some cases, surgeons work together to provide the most effective care [1]. Tennis elbow affects 1 - 3% of general population. Lateral epicondylitis can occur during activities that require repeated supination and pronation of the forearm with the elbow in near full extension. Most of patients with tennis elbow can be managed conservatively; treatment is successful in 95% of patients [2]. Pain usually is exacerbated by resisted wrist dorsiflexion and forearm supination and there is pain when grasping objects. Plain radiographs usually are negative occasionally calcific tendinitis may be present [3]. MRI shows tendon thickening with increased T1 and T2 signals. Microscopic findings show immature reparative tissue that resembles angiofibroblastic fibroplasias. We evaluated short term results of local steroid injections in the management of tennis elbow.

2. Materials and Methods

The present study is the effectiveness of local corticosteroid injection in the management of tennis elbow" was conducted in postgraduate Department of Orthopaedics, Sree Mookambika institute of medical sciences from September 2020 to August 2021.

60 patients with tennis elbow of either sex were included in the study. The patients with more than 3 months of complaints duration and those recently treated with corticosteroids or physiotherapy were excluded. Also patients having bilateral elbow involvement and those having other elbow pathologies were excluded from this study. A record of patient's pain using visual analogue scale (VAS) was obtained at the start of study using a range of 0 to 10, with 0 representing no pain and 10 worst pains ever experienced. After taking written informed consent all the patients received local steroid injection (Methylprednisolone 40 mg mixed with 1 ml injection of lignocaine 2%) Patients were followed at six weeks and final follow-up was done at 3 months.

3. Results

The patients in our study ranged in age from 20 to 60 years with mean age of 44 years. 52 (65%) patients were males and 28 (35%) patients were females. Right side was involved in 48 (57%) patients and left side was involved in 32 (40%) patients, right side was involved more than left side. In our patients, the median VAS pain score at the start of study was 7 (range 4-10) which decreased to a mean score of 2 (range 0-8) at 6 weeks' time. Statistical analysis revealed a significant decrease in score ($p < .001$). But at 3 months (12 weeks) follow majority of patients had recurrence of symptoms VAS score at 3 months ranged from 0 to 10 with median VAS score of 6.



Fig 1: Treatment of tennis elbow

Out of 60 patients in our study 41 (68.3%) patients obtained complete relief of pain at six weeks' time. But at final follow-up of 12 weeks only 14 (24.35 %) patients were pain free. At 12 weeks follow-up 40% (24) of patients had recurrence of symptoms.

Table 1: VAS score and duration

Visual Analogue Scale (VAS)	At the start of study	At 6 weeks	At 12 weeks
0 - 3	0	42 (70.00%)	14 (24.35 %)
4 - 7	20 (31.75 %)	11 (17.50 %)	22 (35.65%)
8 - 10	40 (68.25 %)	7 (12.50 %)	24 (40.00%)

4. Discussion

Tennis Elbow is the most common overuse syndrome in the elbow. It is a tendinopathy injury involving the extensor muscles of the forearm. Tennis elbow is a common clinical disorder of unknown etiology.

- Haker *et al.* in his study gave a comparison of local corticosteroid injection with epicondylitis bandage and splinting [4]. The results for steroid injection was found to be better in two weeks time, but recurrence was detected in 44% of patients in 6 months and results of physical examination were almost similar in both groups at 12 months time [7].
- Smidt *et al.* in his studies reported that corticosteroid injection was more effective in 3-6 months time compared to control or drugs group but at 3-12 months the results of injections was not better than control [5].
- Bisset *et al.* reported that the local corticosteroid injections are effective in short term, but he also stated that results were worst as compared to the other treatment modalities like physiotherapy [6].
- Gosens *et al.* made a comparison between the results of two groups of patients with chronic lateral epicondylitis. The first group was treated by PRP injection, and the second group was treated by corticosteroid injection; both groups significantly improved across time [2]. After 2 years of follow-up, when he checked the DASH score, the score of the corticosteroid group returned to baseline levels while those of the PRP group was drastically improved.

5. Conclusion

In patients with tennis elbow use of methylprednisolone gave good response for short term use and showed to be an excellent treatment. This procedure is not expensive, mini-invasive, with low surgical risk and short hospitalization.

6. Reference

- Campbell's operative orthopaedics; volume three-11th edition, page 19087
- Gosens T, *et al.* Ongoing positive effects of platelet Rich plasma versus corticosteroid injection in lateral epicondylitis: A double Blind Randomized controlled trial: Platelet rich plasma versus corticosteroid injection with two year follow-up. *Am J Sports Med.* 2011;39:1200-12008.
- Lauren Gorelick, Ayal Rozan Gorelick, *et al.* Lateral Epicondylitis injection therapy: A safety and efficacy analysis of Hyaluronate versus corticosteroid injection. *Adv tech Bio/Med.* 2015;3:2
- Haker E, *et al.* Lateral epicondylalgia: A diagnostic and therapeutic challenge. *Karolinska Institute; Stockholm,* 1991.
- Smidt N, Vander Windt D, *et al.* Corticosteroid injections, physiotherapy or a wait and watch policy for lateral epicondylitis: A randomized controlled trial. *Lancet.* 2002;359:657-62.
- Bisset L, *et al.* Mobilization with movement and exercise, corticosteroid injection, or wait and see for tennis elbow: Randomized trail *BMJ.* 2006;333:939.
- Gulzar Saeed Ahmad, Muhammad Ah., *et al.* Tennis elbow: Role of local steroid injection. *J Ayoub Med Coll Abbottbad,* 2012, 24(2).
- Runge W, Word LE, Coonrad R, *et al.* Microscopic histopathology of chronic refractory lateral epicondylitis, *Am J sports Medicine.* 1992;20:746.