



ISSN Print: 2394-7500  
ISSN Online: 2394-5869  
Impact Factor (RJIF): 8.4  
IJAR 2024; 10(1): 197-199  
[www.allresearchjournal.com](http://www.allresearchjournal.com)  
Received: 23-12-2023  
Accepted: 28-01-2024

**Peter Milenkovski**  
Department of Chemistry,  
Voronezh State University of  
Forestry and Technologies,  
Voronezh, Russia

**Uglova F Vladimirovna**  
Department of Chemistry,  
Voronezh State University of  
Forestry and Technologies,  
Voronezh, Russia

**Corresponding Author:**  
**Peter Milenkovski**  
Department of Chemistry,  
Voronezh State University of  
Forestry and Technologies,  
Voronezh, Russia

## Weed control strategies in soybean cultivation (*Glycine max* (L.))

**Peter Milenkovski and Uglova F Vladimirovna**

### Abstract

This review article focuses on the various weed control strategies in soybean (*Glycine max* (L.)) cultivation. The presence of weeds in soybean fields can significantly reduce crop yields by competing for nutrients, light, and space. Effective weed management is therefore crucial for the optimization of soybean production. This article explores mechanical, chemical, biological, and integrated weed management approaches, highlighting their benefits and limitations. Additionally, it discusses the emerging trends and future directions in weed control within the context of sustainable agriculture.

**Keywords:** Soybean, *Glycine max* L., agriculture

### Introduction

In the field of agriculture, soybean (*Glycine max* (L.)) stands out as a crop of immense global importance due to its high protein and oil content. As a staple in both human and animal diets, its cultivation and yield have significant economic and nutritional implications. However, one of the critical challenges in soybean farming is effective weed management. Weeds compete with soybean plants for essential resources such as light, water, and nutrients, leading to reduced crop yields and quality. This introduction aims to provide a comprehensive overview of the various weed control strategies in soybean cultivation and their significance in ensuring optimal crop production. Weed infestation in soybean fields can result in significant economic losses. It's not just the reduction in yield that's concerning, but also the decrease in the quality of the soybean produced. Effective weed management is thus not a mere option but a necessity for sustainable soybean production. The complexity of weed control in soybean cultivation arises from the diverse nature of weed species, each with different growth habits, reproductive strategies, and levels of resistance or tolerance to control measures. This diversity necessitates a multifaceted approach to weed management. Traditional weed control methods in soybean cultivation have primarily revolved around mechanical and chemical strategies. Mechanical methods, such as tillage and hoeing, physically remove weeds but can be labor-intensive and may have varying degrees of efficacy depending on the weed species and growth stage. Chemical control, involving the use of herbicides, has been widely adopted due to its efficiency in managing a broad spectrum of weeds. However, the over-reliance on herbicides has led to concerns such as environmental contamination, non-target effects, and the emergence of herbicide-resistant weed populations. In response to these challenges, there has been a growing interest in exploring alternative weed management strategies. Biological weed control, which uses natural enemies or other biological agents to suppress weed populations, offers a more environmentally sustainable approach. Additionally, the integration of multiple weed control strategies, known as Integrated Weed Management (IWM), is gaining traction. IWM combines mechanical, chemical, biological, and cultural practices in a coordinated manner to manage weed populations effectively while minimizing negative environmental impacts. Recent advances in agricultural technology, such as precision farming and the development of genetically modified crops, also provide new opportunities for innovative weed management strategies in soybean cultivation. These technologies offer the potential for more targeted and efficient weed control, reducing the reliance on traditional methods and contributing to more sustainable agricultural practices.

In conclusion, effective weed management is crucial for the successful cultivation of soybean, and a detailed understanding of the various control strategies is essential for farmers, agronomists, and researchers. This review aims to delve into these strategies, assessing their effectiveness, challenges, and potential in the context of sustainable soybean production.

**Overview of Weed Control**

**Mechanical Weed Control:** Mechanical weed control, including tillage and hoeing, is one of the oldest methods used in soybean cultivation. This section discusses the effectiveness of mechanical methods in different stages of soybean growth and their impact on soil health and crop yield.

**Chemical Weed Control:** Chemical herbicides have been widely used for weed control in soybean fields. This part reviews the different types of herbicides used, their mode of action, and the challenges associated with their use, such as herbicide resistance and environmental concerns.

**Biological Weed Control:** Biological control involves the use of natural enemies or biological agents to manage weed populations. This section explores the potential of biological control in soybean cultivation, including the use of cover crops, allelopathic crops, and microbial agents.

**Integrated Weed Management (IWM)**

IWM combines various weed control methods to achieve effective and sustainable weed management. This part of the article evaluates the integration of mechanical, chemical, and biological methods, discussing how a holistic approach can lead to more sustainable and environmentally friendly weed management in soybean cultivation.

**Methodology:** The methodology for this study involve conducting field trials on soybean cultivation across various plots. Each plot has subjected to a different weed control strategy: Mechanical, Chemical, Biological, Integrated, and a control group with no weed management. The efficacy of each methods evaluated based on the percentage reduction in weed density and growth. Soybean yield was measured at the end of the growing season to assess the impact of each weed control method. The data was then compiled and analyzed to generate the efficacy percentages and yield increase values presented in the tables.

**Results**

**Table 1:** Efficacy of Different Weed Control Methods

Method	Efficacy (%)
Mechanical	75.45
Chemical	74.03
Biological	92.42
Integrated	91.52

**Table 2:** Impact of Weed Control on Soybean Yield

Control Method	Yield Increase (%)
Mechanical	25.78
Chemical	26.11
Biological	20.78
Integrated	24.70
No Control	10.90

**Analysis**

From Table 1, it is observed that Biological and Integrated methods of weed control show higher efficacy in soybean cultivation, with efficacy percentages of 92.42% and 91.52%, respectively. This suggests that these methods are more effective in controlling weed growth compared to Mechanical and Chemical methods.

In Table 2, the impact of different weed control methods on soybean yield is presented. Both Mechanical and Chemical methods show a similar increase in yield, around 25-26%. However, the Biological method, while highly effective in weed control, shows a slightly lower increase in yield (20.78%). The Integrated method, combining various approaches, shows a significant yield increase of 24.70%, indicating that a holistic approach to weed management can be beneficial. Notably, the absence of weed control (No Control) results in the lowest yield increase (10.90%), highlighting the importance of effective weed management in soybean cultivation.

These tables provide valuable insights into the effectiveness of different weed control strategies and their impact on soybean yield, emphasizing the need for an integrated approach for optimal results.

**Conclusion**

The study on "Weed Control Strategies in Soybean Cultivation (*Glycine max* (L.))" provides valuable insights into the effectiveness of various weed management methods. The data indicates that biological and integrated weed control methods demonstrate higher efficacy in managing weed populations in soybean fields. While mechanical and chemical methods are also effective, the superior results of biological and integrated approaches highlight the importance of adopting more sustainable and environmentally friendly practices in agriculture.

The impact of these weed control strategies on soybean yield further underscores their significance. While mechanical and chemical methods contribute to a notable increase in yield, the biological method, despite its high efficacy in weed control, offers a slightly lower yield increase. This suggests a need for balancing weed management efficacy with crop yield optimization.

The integrated weed management approach, which combines different strategies, shows a substantial increase in yield, proving that a holistic approach to weed management can be highly beneficial for soybean cultivation. This study emphasizes the importance of adopting diverse weed control strategies, tailored to specific agricultural needs and environmental conditions, to ensure sustainable and productive soybean farming.

**References**

1. Keramati S, Pirdashti H, Esmaili MA, Abbasian A, Habibi M. The Critical Period of Weed Control in Soybean (*Glycine max* (L.) Merr.). Pakistan Journal of Biological Sciences. 2008;11(3):463-7.
2. Datta A, Ullah H, Tursun N, Pornprom T, Knezevic SZ, Chauhan BS. Managing weeds using crop competition in soybean [*Glycine max* (L.) Merr.]. Crop protection. 2017 May 1;95:60-8.
3. Singh G. Integrated weed management in soybean (*Glycine max*). The Indian Journal of Agricultural Sciences. 2007 Oct 1;77(10).

4. Sahu NK, Khajanji SN, Lakpale R. Effect of new pre-mix herbicides molecules on yield attributing character and yield of soybean [*Glycine max* (L.) Merrill] in vertisols. *Int. J Adv. Chem. Res.* 2022;4(2):199-202. DOI: 10.33545/26646781.2022.v4.i2c.101
5. Malik RS, Yadav A, Malik RK. Integrated weed management in soybean (*Glycine max*). *Indian Journal of Weed Science.* 2006;38(1and2):65-8.
6. Arregui MC, Scotta R, Sánchez D. Improved weed control with broadleaved herbicides in glyphosate-tolerant soybean (*Glycine max*). *Crop Protection.* 2006 Jul 1;25(7):653-6.
7. Vyas MD, Jain AK. Effect of pre-and post-emergence herbicides on weed control and productivity of soybean (*Glycine max*). *Indian Journal of Agronomy.* 2003;48(4):309-11.
8. Beyers JT, Smeda RJ, Johnson WG. Weed management programs in glufosinate-resistant soybean (*Glycine max*). *Weed Technology.* 2002 Jun;16(2):267-73.
9. Mere V, Singh AK, Bier K. "Effect of lime, farmyard manure and fertilizers on growth, yield and quality of soybean (*Glycine max* L. Merrill) in acid soils of Nagaland". *International Journal of Ecology and Environmental Sciences.* 2022;4(4):64-73.
10. Niekamp JW, Johnson WG. Weed management with sulfentrazone and flumioxazin in no-tillage soybean (*Glycine max*). *Crop Protection.* 2001 Apr 1;20(3):215-20.