



ISSN Print: 2394-7500
 ISSN Online: 2394-5869
 Impact Factor (RJIF): 8.4
 IJAR 2024; 10(11): 315-318
www.allresearchjournal.com
 Received: 12-09-2024
 Accepted: 18-10-2024

Dr. Titlee Majumder
 Assistant Professor, Swami
 Vivekananda University,
 Barrackpore, Kolkata,
 West Bengal, India

Importance of excitotoxicity in stress: A review

Dr. Titlee Majumder

DOI: <https://doi.org/10.22271/allresearch.2024.v10.i11e.12178>

Abstract

Excitotoxicity refers to the process where nerve cells are damaged or killed due to excessive stimulation by neurotransmitters, primarily glutamate. This overactivation of receptors, such as NMDA (N-methyl-D-aspartate) receptors, causes an influx of calcium ions into the neurons, leading to cell damage or death. It is a key factor in various neurodegenerative diseases and acute neurological conditions like stroke and traumatic brain injury. Many researchers suggested that chronic stress can exacerbate excitotoxicity by increasing the release of glutamate in the brain. There are numerous factors play extremely important roles in many endorsing the excitotoxicity are increased glutamate levels, vulnerability of the neurones and neuro-degeneration in the diseases like Parkinsons. Several protective measures are obtained by the review of many researches. Effective stress management, including mindfulness, exercise, and therapy, can help regulate glutamate levels and reduce the risk of excitotoxicity. Enhancing antioxidant defences through diet or supplementation may protect neurons from the oxidative stress induced by excitotoxicity. Drugs that block NMDA receptors or regulate glutamate activity are being explored as treatments for conditions associated with excitotoxicity.

Keywords: Excitotoxicity, Parkinson, neuro degeneration, stress, NMDA

Introduction

Excitotoxicity is a process where neurons are damaged and killed by excessive stimulation by neurotransmitters, particularly glutamate. When there's an overabundance of glutamate in the synaptic cleft, it can lead to overactivation of glutamate receptors. This excessive stimulation causes an influx of calcium ions into the neurons, triggering a cascade of events that can result in cell damage and death. Excitotoxicity is implicated in various neurological disorders, such as stroke, Alzheimer's disease, and multiple sclerosis. The phenomenon highlights the delicate balance that the brain maintains between neurotransmission and neuroprotection^[1].

Excitotoxicity can arise from several factors, including^[2]

- 1. Ischemia:** Reduced blood flow (e.g., during a stroke) leads to energy failure, causing cells to release excess glutamate.
- 2. Trauma:** Brain injuries can cause cell damage and disrupt normal neurotransmitter levels, increasing glutamate release.
- 3. Neurodegenerative Diseases:** Conditions like Alzheimer's, Parkinson's, and Huntington's disease often involve chronic excitotoxicity due to neuronal degeneration and imbalanced neurotransmitter signalling.
- 4. Seizures:** Epileptic activity can lead to high levels of glutamate, promoting excitotoxic effects on neurons.
- 5. Toxic Exposure:** Certain substances, such as heavy metals and some drugs, can disrupt glutamate signalling and lead to excitotoxicity.
- 6. Inflammation:** Neuroinflammatory responses can increase the release of glutamate and reduce its reuptake, contributing to excitotoxic conditions.
- 7. Metabolic Disorders:** Conditions like diabetes can alter energy metabolism in neurons, increasing vulnerability to excitotoxic damage.

Literature Review

Multiple researches suggested that there are many reasons well established to make remarkable evidences proving excitotoxicity amongst the individuals, which are as follows^[3]; overstimulation of glutamate receptors, leading to a series of physiological mechanisms

Corresponding Author:
Dr. Titlee Majumder
 Assistant Professor, Swami
 Vivekananda University,
 Barrackpore, Kolkata,
 West Bengal, India

for example; few evidences suggest that, overactivation of glutamate receptors lead to excess glutamate binding to its receptors (primarily NMDA and AMPA receptors), causing prolonged depolarization of the postsynaptic neuron [4]. Other researches suggested that activation of calcium receptors allows excessive calcium ions (Ca^{2+}) to flow into the neuron. Calcium is a crucial signalling molecule, but too much of it can be toxic [5]. Other researches, showed elevated intracellular calcium levels activate various enzymatic pathways, including: (a) Phospholipase A2: This can lead to the release of arachidonic acid, promoting inflammation and further damaging cell membranes. (b) Nitric Oxide Synthase: Excess calcium can activate this enzyme, resulting in increased production of nitric oxide, which can cause oxidative stress [6]. Numerous evidences also show that, high levels of calcium can overwhelm mitochondria, impairing their ability to produce ATP and leading to energy deficits [7]. Mitochondrial dysfunction also increases the production of reactive oxygen species (ROS), which can damage cellular components [8]. Activation of numerous cell death pathways initiate combination of oxidative stress, inflammation, and energy failure can trigger apoptotic (programmed cell death) and necrotic pathways, leading to irreversible neuron damage [9]. Under normal conditions, glial cells (like astrocytes) help recycle glutamate. In excitotoxic scenarios, the ability of these cells to uptake and clear glutamate may be compromised, exacerbating the problem [10]. These mechanisms collectively contribute to neuronal injury and death, highlighting the importance of tightly regulating glutamate levels and receptor activity in the brain [11]. It has been already researched and established that excitotoxicity refers to the process by which excessive activation of glutamate receptors leads to neuronal damage and cell death [12]. This can occur due to a variety of factors, including: Ischemia (lack of blood flow), Traumatic brain injury, Neurodegenerative diseases (like Alzheimer's and Parkinson's), Chronic stress etc. When neurons are exposed to high levels of glutamate, they can become overstimulated, leading to increased intracellular calcium levels, oxidative stress, and ultimately cell death [13]. This process is often associated with inflammation and the activation of apoptotic pathways.

Stress exacerbating excitotoxicity

Whereas, stress can be defined as, particularly chronic stress, can exacerbate excitotoxicity through several mechanisms [14, 15]:

- **Hormonal Changes:** Stress increases levels of cortisol and other stress hormones, which can affect neuronal health and synaptic plasticity.
- **Inflammation:** Chronic stress can lead to increased inflammatory markers, which can further promote excitotoxicity.
- **Glutamate Release:** Stress can alter glutamate dynamics, potentially leading to heightened levels of this neurotransmitter in synaptic clefts.
- **Energy Metabolism:** Stress impacts the brain's energy metabolism, which can compromise neuronal survival during excitotoxic events.

Stress can have complex effects on the body, and while the parasympathetic nervous system (PNS) is generally responsible for promoting relaxation and recovery, chronic stress can lead to dysfunction in this system [16]. Here are

some research-based evidences present in which it has been showed that chronic stress as well as acute stress both give extreme effects upon the central nervous system in all sensory and motor paths [17]. In many researches it has been showed that, the body's stress response activates the sympathetic nervous system (SNS), which can overshadow the PNS. Prolonged activation of the SNS can lead to decreased PNS activity, making it harder to relax and recover [16]. It has a potential role creating Impact on Heart Rate Variability (HRV). A healthy PNS contributes to high HRV, indicating good adaptability and resilience to stress. Chronic stress can lower HRV, suggesting reduced parasympathetic activity and poorer recovery [17]. Many stresses induced abnormalities are evident in many researches that can be manifest as digestive issues (e.g., IBS), sleep disturbances, and chronic pain, often linked to impaired parasympathetic function. Even the body may struggle to maintain homeostasis due to this imbalance [18, 19]. Even the mental health can be compromised due to elevated levels of stress with holding reduced parasympathetic activity which can contribute to anxiety and depression, further perpetuating the cycle of stress and dysfunction. A weakened PNS might hinder emotional regulation and resilience [20]. The lifestyle factors like- poor sleep, unhealthy eating, and lack of physical activity can exacerbate the effects of stress on the Peripheral Nervous System (PNS). Techniques like mindfulness, deep breathing, and regular exercise can help restore balance [21]. Drug & therapy induced evidences are very prominent on this note with multiple approaches like, stress management techniques, such as yoga, meditation, and biofeedback, can enhance PNS function. Professional support from therapists or counsellors can also be beneficial for managing chronic stress [22, 23]. If you're experiencing significant stress, consider exploring relaxation techniques or speaking with a healthcare professional for tailored advice. Excitotoxicity and neurodegeneration are closely related concepts, particularly in the context of neurodegenerative diseases [24]. Here's an overview of each and their interplay: under normal conditions, glutamate is crucial for synaptic transmission and plasticity. Excessive glutamate release or impaired reuptake can lead to overactivation of glutamate receptors [(e.g., *N*-methyl-*D*-aspartic acid (NMDA), α -amino-3-hydroxy-5-methylisoxazole-4-propionate (AMPA)] [25]. This overactivation causes a cascade of intracellular events, including excessive calcium influx, oxidative stress [26].

Excitotoxicity and neurodegenerative disorders

Neurodegenerative disorders are considered to be the heterogenous set of disorders amongst all the population. The distinct disorders show variety of clinical phenotypes and genotypes [26]. The excitotoxicity based neurological disorders are basically based upon the elderly population rather the geriatric subjects. The cases which give emphasis upon the roles of excitatory neurotransmitters within the cerebral cortex and the hippocampal region relating many diseases by induced forgetfulness [27]. The regulatory releases of excitatory amino acids lead to contribute to variety of clinical aspects like schizophrenia, progressive dementia and many other neuropsychiatric syndromes etc. [28]. The excitotoxicity mainly effects upon glutaminergic neurons. Under the basic physiological conditions, the glutamate effects upon altering the activity of the cationic

channels regulating the excitatory response. The hypothesis derived from various researches show that the classical or traditional pathways of glutamic acid eccentric activities include *N*-methyl-*D*-aspartic acid (NMDA), α -amino-3-hydroxy-5-methylisoxazole-4-propionate (AMPA) and kainic acid (KA) receptors [29]. At first the NMDA receptors induce calcium ion activities, in excess of it leads to nerve injuries. Many researches showed that Huntington's disease is a clinical condition elevate the expressions of NMDA receptor followed by voltage calcium ion channels [30]. Even the same cascade-based alterations are observed within the non-human rodents. The increased catabolic activities of calcium ions lead to activate apoptosis followed by necrosis [31]. On the other hand, the AMPA receptor alteration in exhibiting calcium ion activity leads to arise hypoxic-ischemic conditions amongst the individuals. Though the scientific evaluations can be also seen for the Parkinson's and Alzheimer's disease [32, 33].

Conclusion

Many researches based upon stress induced by excitotoxicity have shown many evidences which found instrumental in increasing oxidative stress. Whereas, extensive experimentation may open newer findings in importance of excitotoxicity induced by stress.

References

- Leipnitz G, da Rosa JS, Wajner M. The Role of Excitotoxicity, Oxidative Stress and Bioenergetics Disruption in the Neuropathology of Nonketotic Hyperglycinemia. *Neurotoxicity Research*. 2024;42(4):32.
- Zhang Z, Gao X, Tian Z, Yang E, Huang Y, Liu D, *et al.* Preso enhances mGluR1-mediated excitotoxicity by modulating the phosphorylation of mGluR1-Homer1 complex and facilitating an ER stress after traumatic brain injury. *Cell Death Discovery*. 2024;10(1):153.
- Nicosia N, Giovenzana M, Misztak P, Mingardi J, Musazzi L. Glutamate-Mediated Excitotoxicity in the Pathogenesis and Treatment of Neurodevelopmental and Adult Mental Disorders. *Int J Mol Sci*. 2024;25(12):6521.
- Saidia AR, François F, Casas F, Mechaly I, Venteo S, Veechi JT, *et al.* Oxidative Stress Plays an Important Role in Glutamatergic Excitotoxicity-Induced Cochlear Synaptopathy: Implication for Therapeutic Molecules Screening. *Antioxidants*. 2024;13(2):149.
- Dionysopoulou S, Wikstrom P, Walum E, Georgakis S, Thermos K. Investigation of the Effects of a Novel NOX2 Inhibitor, GLX7013170, against Glutamate Excitotoxicity and Diabetes Insults in the Retina. *Pharmaceuticals*. 2024;17(3):393.
- Khazaal HT, El-Sayed EK, Mansour YE, Ibrahim RR, Bishr M, El Dib RA, Soliman HS. Neuroprotective activity of *Colocasia esculenta* (L.) Schott leaves against monosodium glutamate-induced excitotoxicity in rats: phytochemical and molecular docking study. *Nat Prod Res*. 2024:1-9.
- Woo MS, Mayer C, Binkle-Ladisch L, Sonner JK, Rosenkranz SC, Shaposhnykov A, *et al.* STING orchestrates the neuronal inflammatory stress response in multiple sclerosis. *Cell*. 2024.
- Yang XM, Yu H, Li JX, Li N, Li C, Xu DH, *et al.* Excitotoxic Storms of Ischemic Stroke: A Non-neuronal Perspective. *Mol Neurobiol*. 2024:1-20.
- Zaki-Dizaji M, Abazari MF, Razzaghi H, Shkolnikov I, Christie BR. GRM7 deficiency, from excitotoxicity and neuroinflammation to neurodegeneration: systematic review of GRM7 deficient patients. *Brain Behav Immun Health*. 2024;100808.
- Kumari S, Dhapola R, Sharma P, Nagar P, Medhi B, HariKrishnaReddy D. The impact of cytokines in neuroinflammation-mediated stroke. *Cytokine Growth Factor Rev*. 2024.
- Anaya-Fernández R, Anaya-Prado R, Anaya-Fernandez MM, Guerrero-Palomera MA, Garcia-Ramirez IF, Gonzalez-Martinez D, *et al.* Oxidative Stress in Cerebral Ischemia/Reperfusion Injury. *OBM Neurobiol*. 2024;8(3):1-15.
- Bondy SC. Mitochondrial Dysfunction as the Major Basis of Brain Aging. *Biomolecules*. 2024;14(4):402.
- Angelova PR, Abramov AY. Interplay of mitochondrial calcium signalling and reactive oxygen species production in the brain. *Biochem Soc Trans*. 2024;52(4):1939-1946.
- Egunlusi AO, Malan SF, Palchykov VA, Joubert J. Calcium Modulating Effect of Polycyclic Cages: A Suitable Therapeutic Approach Against Excitotoxic-Induced Neurodegeneration. *Mini Rev Med Chem*. 2024;24(13):1277-1292.
- Ekundayo BE, Obafemi TO, Adewale OB, Obafemi BA, Oyinloye BE, Ekundayo SK. Oxidative Stress, Endoplasmic Reticulum Stress and Apoptosis in the Pathology of Alzheimer's Disease. *Cell Biochem Biophys*. 2024:1-21.
- Ekundayo BE, Obafemi TO, Adewale OB, Obafemi BA, Oyinloye BE, Ekundayo SK. Oxidative Stress, Endoplasmic Reticulum Stress and Apoptosis in the Pathology of Alzheimer's Disease. *Cell Biochem Biophys*; c2024. p. 1-21.
- Egunlusi AO, Joubert J. NMDA Receptor Antagonists: Emerging Insights into Molecular Mechanisms and Clinical Applications in Neurological Disorders. *Pharmaceuticals*. 2024;17(5):639.
- Zakaria N, Fadhlina A, Sheikh H, Hairani M, MohdFauzi M, AbdulMajid F. Stress-Relieving Properties of a Polyherbal Blend with *Syzygium aromaticum* L. and *Coffea canephora* Pierre ex A. Froehner: A Review and Bibliometric Analysis. *World J Biol Psychiatry*. 2024:1-18.
- Castillo-Vazquez SK, Massieu L, Rincón-Heredia R, García-delaTorre P, Quiroz-Baez R, Gomez-Verjan JC, *et al.* Glutamatergic neurotransmission in aging and neurodegenerative diseases: A potential target to improve cognitive impairment in aging. *Arch Med Res*. 2024;55(6):103039.
- Stojanovic M, Rai V, Agrawal DK. Effect of Electromagnetic Field on Proliferation and Migration of Fibroblasts and Keratinocytes: Implications in Wound Healing and Regeneration. *J Biotechnol Biomed*. 2024;7:387-399.
- Tasca CI, Zuccarini M, Di Iorio P, Ciruela F. Lessons from the physiological role of guanosine in neurodegeneration and cancer: Toward a multimodal mechanism of action? *Purinergic Signal*. 2024:1-16.

22. Owjifard M, Rahimian Z, Karimi F, Borhani-Haghighi A, Mallahzadeh A. A comprehensive review on the neuroprotective potential of resveratrol in ischemic stroke. *Heliyon*. 2024.
23. Islam MR, Jony MH, Thufa GK, Akash S, Dhar PS, Rahman MM, *et al.* A clinical study and future prospects for bioactive compounds and semi-synthetic molecules in the therapies for Huntington's disease. *Mol Neurobiol*. 2024;61(3):1237-1270.
24. Chrobak AA, Pańczyszyn-Trzewik P, Król P, Pawelec-Bąk M, Dudek D, Siwek M. New light on prions: putative role of PrPc in pathophysiology of mood disorders. *Int J Mol Sci*. 2024;25(5):2967.
25. Saramak K, Szejko N. Introductory Chapter: Motor Neurons–New Insights. In: *Motor Neurons–New Insights*. IntechOpen; c2024.
26. Yang J, Yin N, Yang R, Faiola F. Role of Pollution-Induced Oxidative Stress and Inflammation in Neurodegenerative Diseases, and the Mechanisms of Traditional Chinese Medicine's Potential Remediation. *Rev Environ Contam Toxicol*. 2024;262(1):9.
27. Richard SA. Elucidating the pivotal molecular mechanisms, therapeutic and neuroprotective effects of lithium in traumatic brain injury. *Brain Behav*. 2024;14(6):e3595.
28. Baburaj R, Sandur VR, Das K. Investigation of the Pro-active Role of Alpha Amyrin Nanoemulsions in Quashing Neurodegeneration, Excitotoxicity, and Neuronal Inflammation-A Combined *in vivo* and *in silico* Approach. *Indian J Pharm Educ Res*. 2024;58(1).
29. Briones-Valdivieso CK, Briones F, Orellana-Urzúa S, Chichiarelli S, Saso L, Rodrigo R. Novel Multi-antioxidant Approach for ischemic stroke therapy targeting the role of oxidative stress. *Biomedicines*. 2024;12(3):501.
30. Oshchepkov DY, Makovka YV, Fedoseeva LA, Seryapina AA, Markel AL, Redina OE. Effect of Short-Term Restraint Stress on the Hypothalamic Transcriptome Profiles of Rats with Inherited Stress-Induced Arterial Hypertension (ISIAH) and Normotensive Wistar Albino Glaxo (WAG) Rats. *Int J Mol Sci*. 2024;25(12):6680.
31. Fu C, Lei Y, Liang L, Jiang J, Qin Y, Lao Y, *et al.* Characterization of HSP90 expression and function following CNS injury. *Neurosci Lett*. 2024;137875.
32. Woo MS, Engler JB, Friese MA. The neuropathobiology of multiple sclerosis. *Nat Rev Neurosci*. 2024:1-21.
33. Theus MH. Neuroinflammation and acquired traumatic CNS injury: a mini review. *Front Neurol*. 2024;15:1334847.